



Product Information Sheet

Monoclonal Anti- β -Amyloid Protein (Sephacrose Bead Conjugate)

Catalogue No. MA1109-S

Immunogen

Synthetic β -amyloid peptide (1-40), conjugated to KLH.

Lot No. 08A12

Purification

Purified by the goat anti-mouse IgG affinity chromatography.

Clone: AP-5

Ig type: mouse IgG1

Formulation

50% slurry in PBS pH 7.2 with 0.01mg NaN₃ preservative.

Size: 200 μ l

Storage

Store at 4°C for frequent use.

Specificity

Human.

No cross reactivity with other proteins.

Description:

This Antagene antibody is immobilized via covalent binding of primary amino groups to N-hydroxysuccinimide (NHS)-activated sephacrose beads. It is useful for immunoprecipitation assays

Recommended application

Immunoprecipitation(IP)

BACKGROUND

Beta amyloid protein is 4.2-kD polypeptide. Glenner and Wong (1984) purified a protein derived from the twisted beta-pleated sheet fibrils present in cerebrovascular amyloidoses and in the amyloid plaques associated with Alzheimer disease. The proteins from both disorders have an identical 28-amino acid sequence. memory deficits in middle-aged Tg2576 mice are caused by the extracellular accumulation of a 56-kDa soluble amyloid-beta assembly. Amyloid beta protein (A β) deposition in the brain is a hallmark of Alzheimer's disease (AD).

REFERENCE

1. Lesne, S.; Koh, M. T.; Kotilinek, L.; Kaye, R.; Glabe, C. G.; Yang, A.; Gallagher, M.; Ashe, K. H. : A specific amyloid-beta protein assembly in the brain impairs memory. *Nature* 440: 352-357, 2006.
2. Lorenzo, A.; Yuan, M.; Zhang, Z.; Paganetti, P. A.; Sturchler-Pierrat, C.; Staufenbiel, M.; Mautino, J.; Vigo, F. S.; Sommer, B.; Yankner, B. A. : Amyloid beta interacts with the amyloid precursor protein: a potential toxic mechanism in Alzheimer's disease. *Nature Neurosci.* 3: 460-464, 2000.

For Research Use Only not for diagnostic and clinical use

Contact: Antagene, Inc. | Tel: 1 (866) 964-2589 | Fax: 1 (888) 225-1868 | Email: Info@antageneinc.com