

Mouse Anti Human CD19

Code: [USC] 19P-HIB19-50

SIZE: 50 TEST

Clone: HIB19

Label•Form: PE conjugation

WSNo: V CD19.11

Isotype: IgG1

Description:

CD19 McAb recognizes a 95 KD type I transmembrane glycoprotein which is restricted B cell antigen. CD19 antigen is expressed on normal and neoplastic B cells and also in some the bone marrow cells. CD19 expression by B progenitor cells is presumably at late pro-B or early pre-B stages around the time of Ig heavy chain rearrangement. Expression persists during all stages of B cell maturation and is lost on terminal differentiation to plasma cells. CD19 antigen is also found on the follicular dendritic cells and the early cells of myelomonocytic lineage but not on normal T cells, NK cells, monocytes, granulocytes, erythrocytes and platelets. In normal peripheral blood, 8-20% of lymphocytes express CD19 antigen. CD19 antigen plays a role in regulating B cell proliferation.

Preparation:

The monoclonal antibody is purified from ascites by protein G affinity chromatography and is conjugated with FITC, R-PE, PE-Cy5 under optimum conditions.

Usage:

The purified reagent is effective for indirect immunofluorescence staining of human cells for flow cytometric analysis and is tested for immunohistochemical staining of acetone-fixed frozen sections. The conjugated reagent (FITC, R-PE) is tested for flow cytometric analysis using 20 μ l/10⁶ cells or 100 μ l peripheral blood cells. The conjugated reagent (PE-Cy5) is tested for flow cytometric analysis using 10 μ l/10⁶ cells or 100 μ l peripheral blood cells.

Storage:

For purified forms, long term storage at -20°C. For conjugated forms, storage at 4°C, should not be frozen and avoid prolonged exposure to light.

References:

Schlossman, S., L.Bloumsell, W.Gilks, et al., eds. 1995. Leucocyte Typing V: White Cell Differentiation Antigens. P:491, 507, 2004 Oxford University Press, New York.



コスモ・バイオ株式会社

〒135-0016 東京都江東区東陽2-2-20 東陽駅前ビル
TEL : (03) 5632-9610
(03) 5632-9620
FAX : (03) 5632-9619
e-mail : mail@cosmobio.co.jp
URL : <http://www.cosmobio.co.jp/>



COSMO BIO CO., LTD.

TOYO EKIMAE BLDG. 2-20, TOYO 2-CHOME,
KOTO-KU. TOKYO 135-0016, JAPAN
TEL : (81)3-5632-9617 FAX : (81)3-5632-9618
e-mail : export@cosmobio.co.jp
URL : <http://www.cosmobio.co.jp/>