

## Mouse Anti Human CD15 Monoclonal Antibody

DMABT-45615MH Mouse(CD15)  
Lot. No. (See product label)

### PRODUCT INFORMATION

<b>Product Overview</b>	Mouse Anti Human CD15
<b>Host</b>	Mouse
<b>Isotype</b>	IgM
<b>Species</b>	Human
<b>Clone</b>	Cv39
<b>Conjugation</b>	N/A
<b>Applications</b>	IHC, ELISA, FCM, IP, WB
<b>Dilution</b>	FCM: 1/100 - 1/200

### PACKAGING

<b>Format</b>	Purified IgM
<b>Protein Concentration</b>	Ig concentration 1.0mg/ml
<b>Buffer</b>	Phosphate buffered saline
<b>Storage</b>	Store at +4 °C or at -20 °C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
<b>Preservative</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Shelf Life</b>	18 months from date of despatch.

### BACKGROUND

<b>Introduction</b>	<p>CD15 (3-fucosyl-N-acetyl-lactosamine) is a cluster of differentiation antigen - an immunologically significant molecule. CD15 is a carbohydrate adhesion molecule (not a protein) that can be expressed on glycoproteins, glycolipids and proteoglycans. CD15 is expressed on Reed-Sternberg cells of Hodgkin's disease and by various other cell types including myeloid cells and epithelial cells. Antibodies to CD15 recognize a pentasaccharide sequence occurring in lacto-N-fucopentaose III ceramide (also referred to as X hapten of Lex) found in higher glycolipids and glycoproteins. A review by Arber et al. has reported that antibodies to CD15 demonstrate positive staining in 87% of Hodgkin's disease including nodular sclerosing, mixed cellularity, and lymphocyte depletion, whereas the lymphocyte predominant variant exhibits a lower rate of positivity (37%). Among non-Hodgkin's lymphoma, 13% express CD15 including 4.1% B-cell, 21% T-cell, and 17% null-cell. CD15 expression has also been demonstrated in acute myeloid leukemia (65%) and chronic myelogenous leukemia (96% chronic phase and 54% blast phase). A relatively low level of CD15 expression has been reported in acute lymphoblastic leukemia (5.7% overall) with positivity observed in 7.7% common or precursor B-cell, 0% B-cell, 7.7% T-cell and 17.3% nullcell. Carcinomas derived from various organs have also been shown to be CD15 positive (56%) including adenocarcinomas, squamous cell carcinomas and undifferentiated large and small cell carcinomas.</p>
---------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

**Keywords**

fucosyltransferase; Alphafucosyltransferase; Alpha 13 fucosyltransferase FucT; Alpha-1 ELAM ligand fucosyltransferase; ELAM-1 ligand fucosyltransferase; ELAM1 ligand fucosyltransferase; ELFT; FCT3A; Fuc-TIV; Fucosyltransferase 4 alpha 1 3 fucosyltransferase myeloid specific; Fucosyltransferase 4; Fucosyltransferase IV; FucT IV; FucT-IV; FUCTIV; FUT4; FUT4\_HUMAN; Galactoside 3 L fucosyltransferase; Galactoside 3-L-fucosyltransferase; LeX; SSEA 1; SSEA1; Stage specific embryonic antigen 1