

Monoclonal Anti-human Neuropilin-1/CD304

Product reference: DDX0440

Description

We have generated a mAb that identifies human pDCs *ex vivo*. In blood, 211H6 antigen is highly expressed on IL3Rα⁺ BDCA-2⁺ CD11c⁻ pDCs. It does not stain lymphocytes, monocytes, basophils and granulocytes. *In vitro* it is also expressed on monocyte-derived DCs. Upon maturation, in IL-3 and CD40L, or with *influenzae* virus, 211H6 is maintained at the cell surface. *In situ* 211H6 recognizes some isolated cells in T cell zones of tonsils, lymphatic vessels and high endothelial venules. The 211H6 antigen (110KDa) has been identified as BDCA-4. Neuropilin-1 is a membrane-bound co-receptor to a tyrosine kinase receptor for both vascular endothelial growth factor (VEGF) and semaphorin (SEMA3A) family members. Neuropilin 1 plays versatile roles in angiogenesis, axon guidance, cell survival, migration and invasion. (*Dzionek A and al.*, 2000 *J. Immunol.*, 165: 6037-6046; *Dzionek A. et al.*, 2002, *Human Immunology*, 63,1133-48).

Clone: 211H6.01 Species: mouse

Specificity: human neuropilin-1 (epitope in extracellular domain)

Immunogen: CD34⁺ derived DCs

Species cross- reactivity: nd IgG1


Formulation/size: Purified: 100 μg in 200μ1 / 50 μg in 100 μl Tris-NaCl pH 8

Coupled: 100 µg in 200µl / 50 µg in 100 µl PBS 50% glycerol

Available formats:

| ueu- | | | |
|----------------|-----------------|---------------------|-------------------------|
| Reference N° | | | |
| 50 μg | 100 µg | Format | Application tested |
| DDX0440P-50 | DDX0440P-100 | Purified | Flow cytometry, IHC, IP |
| DDX0440A488-50 | DDX0440A488-100 | Alexa-fluor® 488 | Flow cytometry |
| DDX0440A546-50 | DDX0440A546-100 | Alexa- fluor® 546 | Flow cytometry |
| DDX0440A647-50 | DDX0440A647-100 | Alexa- fluor® 647 | Flow cytometry |
| DDX0440B-50 | DDX0440B-100 | Biotin (on request) | Flow cytometry, IHC, IP |

Applications tested: Flow cytometry, IHC



Usage recommendation: *This monoclonal antibody may be used between 5-20 µg/ml

*Optimal dilution should be determined by each laboratory for each

application

*Coupled antibody: to maintain RT before using

Aliquot storage conditions: -20°C. KEEP CONTENTS STERILE: no preservative.

<u>Purified</u> antibodies: avoid repeated freeze/thaw cycles. <u>Coupled</u> antibodies: glycerol protects from freezing.