

RPA371Bo01 50µg
Recombinant Interferon Gamma Induced Protein 10kDa (IP10)
Organism Species: Bos taurus; Bovine (Cattle)
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[**PROPERTIES**]

Residues: Val20~Ala102

Tags: Two N-terminal Tags, His-tag and T7-tag

Accession: Q2KIQ8

Host: *E. coli*

Subcellular Location: Secreted.

Purity: >95%

Endotoxin Level: <1.0EU per 1µg
(determined by the LAL method).


Formulation: Supplied as lyophilized form in 20mM Tris,
150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT,
0.01% sarcosyl, 5% trehalose, and preservative.

Predicted isoelectric point: 10.5

Predicted Molecular Mass: 13.2kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)



[**USAGE**]

Reconstitute in sterile ddH₂O.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

V PLSRNTRCSC IEISNGSVNP RSLEKLEVIP ASQSCRVEI IATMKKNGEK RCLNPESKTI
KNLLKAINKQ RTKRSPRTRK EA

[REFERENCES]

1. Hudgens T., *et al.* (2007) Submitted to the EMBL/GenBank/DDBJ databases.
2. Luster AD., *et al.* (1987) Proc. Natl. Acad. Sci. U.S.A. 84 (9): 2868–71.
3. Angiolillo AL., *et al.* (1995) J. Exp. Med. 182 (1): 155–62.
4. Booth V., *et al.* (2002) Biochemistry 41 (33): 10418–25.