



P91054Mu01 100µg **Apoptotic Peptidase Activating Factor 1 (APAF1)** Organism: Mus musculus (Mouse) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

6th Edition (Revised in March, 2013)

kDal

94 66.2

45

33

26

[PROPERTIES]

Residues: Arg872~Phe1135 (Accession # O88879),

with N-terminal His-Tag.

Host: E. coli

Subcellular Location: Cytoplasm.

Purity: >95%

Endotoxin Level: <1.0EU per 1µg

Formulation: Supplied as lyophilized form in PBS,

Predicted isoelectric point: 6.3

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

20 (determined by the LAL method). 14.4 pH7.4, containing 5% sucrose, 0.01% sarcosyl. 15% SDS-PAGE Predicted Molecular Mass: 30.9kDa

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.





[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with N-terminal His-Tag, its sequence is listed below. MGHHHHHHSGSEF- RLKVADCRG HLSWVHGVMF SPDGSSFLTA SDDQTIRVWE TKKVCKNSAI VLKQEIDVVF QENETMVLAV DNIRGLQLIA GKTGQIDYLP EAQVSCCCLS PHLEYVAFGD EDGAIKIIEL PNNRVFSSGV GHKKAVRHIQ FTADGKTLIS SSEDSVIQVW NWQTGDYVFL QAHQETVKDF RLLQDSRLLS WSFDGTVKVW NVITGRIERD FTCHQGTVLS CAISSDATKF SSTSADKTAK IWSFDLLSPL HELKGHNGCV RCSAF