

APC128Ra01 200µg

Active Gremlin 1 (GREM1)

Organism Species: Rattus norvegicus (Rat)

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

1st Edition (Apr, 2016)

[PROPERTIES]

Source: Prokaryotic expression.

Host: *E. coli*

Residues: Lys25~Asp184

Tags: N-terminal His-tag

Purity: >95%

Buffer Formulation: 100mM NaHCO₃, 500mM NaCl, pH8.3, containing 1mM DTT, 0.01% sarcosyl, 5%Trehalose.

Applications: Cell culture; Activity Assays.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 9.6

Predicted Molecular Mass: 22.0kDa

Accurate Molecular Mass: 24kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 100mM NaHCO₃, 500mM NaCl (pH8.3) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

```
KKKGSQ GAIPPPDKAQ HNDSEQTQSP  
PQPGSRTRGR GQGRGTAMPG EEVLESSQEA LHVTERKYLK RDWCKTQPLK  
QTIHEEGCNS RTIINRFCYG QCNSFYIPRH IRKEEGSFQS CSFCKPKKFT  
TMMVTLNCP E LQPPTKKKRV TRVKQCR CIS IDLD
```

[ACTIVITY]

Gremlin (GREM1) is an inhibitor in the TGF beta signaling pathway. GREM1 and other BMP antagonists are important in the survival of cancer stroma survival and proliferation in some cancers. The protein expression is found in many cancers and is thought to play important roles in uterine cervix, lung, ovary, kidney, breast, colon, pancreas, and sarcoma carcinomas. A proliferation assay was conducted to detect the bioactivity of recombinant rat GREM1 using A549 cells. Briefly, A549 cells were seeded into triplicate wells of 96-well plates at a density of 5,000 cells/well and allowed to attach, replaced with serum-free overnight, then the medium was replaced with 2% serum standard DMEM prior to the addition of various concentrations of GREM1. After incubated for 96h, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10 μ L of CCK-8 solution was added to each well of the plate, then the absorbance at 450nm was measured using a microplate reader after incubating the plate for 1-4 hours at 37°C. Proliferation of A549 cells after incubation with GREM1 for 96h observed by inverted microscope was shown in Figure1. Cell viability was assessed by CCK-8 (Cell Counting Kit-8) assay after incubation with recombinant GREM1 for 96h. The result was shown in Figure 2. It was obvious that GREM1 significantly increased cell viability of A549 cells.




Figure 1. Cell proliferation of A549 cells after stimulated with GREM1.
(A) A549 cells cultured in DMEM, stimulated with 0.001ng/mL GREM1 for 96h;
(B) Unstimulated A549 cells cultured in DMEM for 96h.




Figure 2. Cell proliferation of A549 cells after stimulated with GREM1.

[IDENTIFICATION]




Figure 3. SDS-PAGE

Sample: Active recombinant GREM1, Rat




Figure 4. Western Blot

Sample: Recombinant GREM1, Rat;

Antibody: Rabbit Anti-Rat GREM1 Ab (PAC128Ra01)

[IMPORTANT NOTE]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.