

MAB995Hu23

**Monoclonal Antibody to Hemojuvelin (HJV)** 

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



### [PROPERTIES]

**Source:** Monoclonal antibody preparation

Host: Mouse

Antibody isotype: IgG1 Kappa

**Purification:** Protein A + Protein G affinity chromatography

Clone number: A7

Traits: Liquid

Concentration: 1mg/ml

**UOM:** 200µl

**Cross Reactivity:** Rat

Applications: WB; IHC; ICC; IP.

[ IMMUNOGEN ]

Immunogen: Recombinant HJV (Leu29~Ala401) expressed in E.coli

Accession No.: RPB995Hu03

[ APPLICATIONS ]

Western blotting: 0.5-3µg/mL;

Immunohistochemistry: 5-30µg/mL;

Immunocytochemistry: 5-30µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300,

50% glycerol.

### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined

# Coud-Clone Corp.

by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

### [ IDENTIFICATION ]






Figure. Western Blot; Sample: Recombinant HJV, Human.



Western Blot; Sample: Rat Heart lysate;
Primary Ab: 2µg/ml Mouse Anti-Human
HJV Antibody Second Ab:
0.2µg/mL HRP-Linked Caprine AntiMouse IgG Polyclonal Antibody
(Catalog: SAA544Mu19)



DAB staining on IHC-P; Samples:
Human Stomach cancer Tissue;
Primary Ab: 30µg/ml Mouse AntiHuman HJV Antibody Second Ab:
2µg/mL HRP-Linked Caprine AntiMouse IgG Polyclonal Antibody
(Catalog: SAA544Mu19)

## [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.