

PAA336Hu01

Polyclonal Antibody to GATA Binding Protein 1 (GATA1)

Organism Species: Homo sapiens (Human)

Instruction manual

←28kDa

9th Edition (Revised in Jul. 2013)

Western Blot

44

33

26

22

18

14

10

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

[PRODUCT INFORMATION]

Immunogen: GATA1, Human

Clonality: Polyclonal

Host: Rabbit

Immunoglobulin Type: IgG

Purification: Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

UOM: 100μg

Sample: Recombinant GATA1, Human

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant GATA1 (Pro150~Gly384) expressed in E.coli.

Accession No.: RPA336Hu01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and

T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF- P ALPSSLPVPN SAYGGPDFSS
TFFSPTGSPL NSAAYSSPKL RGTLPLPPCE ARECVNCGAT ATPLWRRDRT GHYLCNACGL
YHKMNGQNRP LIRPKKRLIV SKRAGTQCTN CQTTTTTLWR RNASGDPVCN ACGLYYKLHQ
VNRPLTMRKD GIQTRNRKAS GKGKKKRGSS LGGTGAAEGP AGGFMVVAGG
SGSGNCGEVA SGLTLGPPGT AHLYQGLGPV VLSGPVSHLM PFPG



[ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against GATA1. It has been selected for its ability to recognize GATA1 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant GATA1 (Pro150~Gly384) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.