



Abbexa Ltd, Innovation Centre, Cambridge Science Park, Cambridge, CB4 0EY, UK
 Telephone: +44 (0) 1223 755950 - Fax: +44 (0) 1223 755951 - E-Mail: info@abbexa.com

Nucleolar Transcription Factor 1 (UBTF) Antibody

Catalogue No.: abx012712



Western blot analysis of extracts from HeLa cells, treated with calyculinA (50ng/ml, 30mins), using UBF (Phospho-Ser484) antibody.



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue using UBF (Phospho-Ser484) antibody.

Rabbit polyclonal antibody against UBF protein. Immunogen region is Internal. Specificity is as follows for the reactive species: H:S484, M:S484, R:S484.

Target: UBF

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

Tested Applications: ELISA, WB, IHC

Recommended dilutions: WB: 1/500 - 1/3000, IHC: 1/50 - 1/100, ELISA: 1/1000. Optimal dilutions/concentrations should be determined by the end user.

Immunogen: The antiserum was produced against synthesized phosphopeptide derived from human UBF around the phosphorylation site of serine 484 (P-E-SP-P-K).

Purification: Purified from rabbit antiserum by affinity chromatography using epitope-specific immunogen.

Form: Liquid

Isotype: IgG

Abbexa Ltd, Innovation Centre, Cambridge Science Park, Cambridge, CB4 0EY, UK
Telephone: +44 (0) 1223 755950 - Fax: +44 (0) 1223 755951 - E-Mail: info@abbexa.com

Conjugation: Unconjugated

Storage: Aliquot and store at -20 °C. Avoid repeated freeze/thaw cycles.

Swiss Prot: [P17480](#)

Buffer: PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl, 0.02% sodium azide, 50% glycerol.

Note: This product is for research use only.