

**Distemper virus Fusion glycoprotein F0 Antibody (FITC)**

Catalogue No.: abx107664

Distemper virus Fusion glycoprotein F0 Antibody (FITC) is a Rabbit Polyclonal antibody against Distemper virus Fusion glycoprotein F0 conjugated to FITC. Under the current model, the protein has at least 3 conformational states: pre-fusion native state, pre-hairpin intermediate state, and post-fusion hairpin state. During viral and plasma cell membrane fusion, the heptad repeat (HR) regions assume a trimer-of-hairpins structure, positioning the fusion peptide in close proximity to the C-terminal region of the ectodomain. The formation of this structure appears to drive apposition and subsequent fusion of viral and plasma cell membranes. Directs fusion of viral and cellular membranes leading to delivery of the nucleocapsid into the cytoplasm. This fusion is pH independent and occurs directly at the outer cell membrane. The trimer of F1-F2 (F protein) probably interacts with H at the virion surface. Upon HN binding to its cellular receptor, the hydrophobic fusion peptide is unmasked and interacts with the cellular membrane, inducing the fusion between cell and virion membranes. Later in infection, F proteins expressed at the plasma membrane of infected cells could mediate fusion with adjacent cells to form syncytia, a cytopathic effect that could lead to tissue necrosis.

**Target:** Distemper virus Fusion glycoprotein F0**Reactivity:** Virus**Host:** Rabbit**Clonality:** Polyclonal**Tested Applications:** ELISA**Recommended dilutions:** Optimal dilutions/concentrations should be determined by the end user.**Immunogen:** Recombinant Canine distemper virus Fusion glycoprotein F0 protein (136-608AA).**Purity:** > 95%**Purification:** Protein G.**Form:** Liquid**Isotype:** IgG**Conjugation:** FITC**Storage:** Aliquot and store at -20 °C. Avoid exposure to light. Avoid repeated freeze/thaw cycles.**Swiss Prot:** [P12569](#)

Abbexa Ltd, Innovation Centre, Cambridge Science Park, Cambridge, CB4 0EY, UK  
Telephone: +44 (0) 1223 755950 - Fax: +44 (0) 1223 755951 - E-Mail: [info@abbexa.com](mailto:info@abbexa.com)

**Gene Symbol:** F

**Buffer:** 0.01 M PBS, pH 7.4, 0.03% Proclin-300 and 50% Glycerol.

**Note:** This product is for research use only.