



Fibroblast Activation Protein Alpha (FAPa) Antibody


Catalogue No.: abx172378




IHC-P analysis of Human Liver Tissue, using Mouse Anti-Human FAPa Antibody (20 µg/ml) and HRP-conjugated Goat Anti-Mouse antibody ([abx400001](#), 2 µg/ml).



IHC-P analysis of Human Small Intestine Tissue, using Mouse Anti-Human FAPa Antibody (20 µg/ml) and HRP-conjugated Goat Anti-Mouse antibody ([abx400001](#), 2 µg/ml).



IHC-P analysis of Human Kidney Tissue, using Mouse Anti-Human FAPa Antibody (20 µg/ml) and HRP-conjugated Goat Anti-Mouse antibody ([abx400001](#), 2 µg/ml).



IHC-P analysis of Human Cerebrum Tissue, using Mouse Anti-Human FAPa Antibody (20 µg/ml) and HRP-conjugated Goat Anti-Mouse antibody ([abx400001](#), 2 µg/ml).

Fibroblast Activation Protein Alpha (FAPa) Antibody is a Mouse Monoclonal antibody against Fibroblast Activation Protein Alpha (FAPa).

Target: Fibroblast Activation Protein Alpha (FAPa)

Reactivity: Human

Abbexa Ltd, Innovation Centre, Cambridge Science Park, Cambridge, CB4 0EY, UK
Telephone: +44 (0) 1223 755950 - Fax: +44 (0) 1223 755951 - E-Mail: info@abbexa.com

Host: Mouse

Clonality: Monoclonal

Tested Applications: WB, IHC, IF/ICC, IP

Recommended dilutions: WB: 0.5-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.

Immunogen: [abx167900](#) - Recombinant FAPa (Ile523-Asp760) expressed in E. coli

Purification: Purified by Protein A and Protein G affinity chromatography.

Form: Liquid

Conjugation: Unconjugated

Storage: Store at 4 °C for frequent use. Aliquot and store at -20 °C for up to one year. Avoid repeated freeze/thaw cycles.

Buffer: PBS, pH 7.4, containing 0.02% NaN₃, 50% glycerol.

Note: This product is for research use only.