

Anti-Human IGF-BP3 Antibody

Catalog # ABG10176

Specification

Anti-Human IGF-BP3 Antibody - Product Information

Application	WB, IHC, E
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

Anti-Human IGF-BP3 Antibody - Additional Information

Preparation

Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hIGF-BP3. Anti-Human IGF-BP3 specific antibody was purified by affinity chromatography employing immobilized hIGF-BP3 matrix.

WesternBlot

To detect hIGF-BP3 by Western Blot analysis this antibody can be used at a concentration of $0.1 - 0.2 \mu g/ml$. Used in conjunction with compatible secondary reagents the detection limit for recombinant hIGF-BP3 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect hIGF-BP3 by sandwich ELISA (using 100 μ l/well antibody solution) a concentration of 0.5 - 2.0 μ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Human IGF-BP3 (60-190BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hIGF-BP3.

Immunohistochemistry

This antibody stained formalin-fixed, paraffin-embedded sections of normal human placenta. The recommended concentrations are 2.5 μ g/ml-0.5 μ g/ml for two hours at room temperature. An HRP-labeled polymer detection system was used with DAB chromogen. Heat induced



antigen retrieval was performed with a pH 6.0 Sodium Citrate buffer. Optimal concentrations and conditions may vary.

Formulation A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage -20°C

Precautions Anti-Human IGF-BP3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Human IGF-BP3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- <u>Flow Cytomety</u>
- <u>Cell Culture</u>