

## Anti-PSMA2 Picoband Antibody Catalog # ABO11706

### Specification

#### Anti-PSMA2 Picoband Antibody - Product Information

Application	WB
Primary Accession	<a href="#">P25787</a>
Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Format	Lyophilized

#### Description

Rabbit IgG polyclonal antibody for Proteasome subunit alpha type-2(PSMA2) detection. Tested with WB in Human;Rat.

#### Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

#### Anti-PSMA2 Picoband Antibody - Additional Information

Gene ID 5683

#### Other Names

Proteasome subunit alpha type-2, 3.4.25.1, Macropain subunit C3, Multicatalytic endopeptidase complex subunit C3, Proteasome component C3, PSMA2, HC3, PSC3

#### Calculated MW

25899 MW KDa

#### Application Details

Western blot, 0.1-0.5 µg/ml, Human, Rat

#### Subcellular Localization


Cytoplasm. Nucleus. Cytoplasm, P-body . Colocalizes with TRIM5 in the cytoplasmic bodies. .

#### Protein Name

Proteasome subunit alpha type-2

#### Contents

Each vial contains 5mg BSA, 0.9mg NaCl,



Western blot analysis of PSMA2 expression in rat testis extract (lane 1) and MCF-7 whole cell lysates (lane 2). PSMA2 at 26KD was detected using rabbit anti- PSMA2 Antigen Affinity purified polyclonal antibody (Catalog # ABO11706) at 0.5 µg/mL. The blot was developed using chemiluminescence (ECL) method .

#### Anti-PSMA2 Picoband Antibody - Background

Proteasome subunit alpha type-2 is a protein that in humans is encoded by the PSMA2 gene. The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit.

0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg NaN<sub>3</sub>.

### **Immunogen**

A synthetic peptide corresponding to a sequence in the middle region of human PSMA2 (82-123aa DYRVLVHRARKLAQQYYLV YQEPIPTAQLVQRVASVMQEYT Q), identical to the related mouse and rat sequences.

### **Purification**

Immunogen affinity purified.

### **Cross Reactivity**

No cross reactivity with other proteins.

### **Storage**

**At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.**

### **Anti-PSMA2 Picoband Antibody - Protein Information**

**Name** PSMA2

**Synonyms** HC3, PSC3

### **Function**

Component of the 20S core proteasome complex involved in the proteolytic degradation of most intracellular proteins. This complex plays numerous essential roles within the cell by associating with different regulatory particles. Associated with two 19S regulatory particles, forms the 26S proteasome and thus participates in the ATP- dependent degradation of ubiquitinated proteins. The 26S proteasome plays a key role in the maintenance of protein homeostasis by removing misfolded or damaged proteins that could impair cellular functions, and by removing proteins whose functions are no longer required. Associated with the PA200 or PA28, the 20S proteasome mediates ubiquitin-independent protein degradation. This type of proteolysis is required in several pathways including spermatogenesis (20S-PA200 complex) or generation of a subset of MHC class I-presented antigenic

peptides (20S-PA28 complex).

**Cellular Location**

Cytoplasm. Nucleus. Note=Colocalizes with TRIM5 in cytoplasmic bodies.  
{ECO:0000250|UniProtKB:P49722}

**Anti-PSMA2 Picoband Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)