

Anti-Rab9 Picoband Antibody
Catalog # ABO12570

Specification

Anti-Rab9 Picoband Antibody - Product Information

Application **WB**
 Primary Accession [P51151](#)
 Host **Rabbit**
 Reactivity **Human, Mouse, Rat**
 Clonality **Polyclonal**
 Format **Lyophilized**

Description

Rabbit IgG polyclonal antibody for Ras-related protein Rab-9A(RAB9A) detection. Tested with WB in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-Rab9 Picoband Antibody - Additional Information

Gene ID 9367

Other Names

Ras-related protein Rab-9A, RAB9A, RAB9

Calculated MW

22838 MW KDa


Application Details

Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat

Subcellular Localization

Cell membrane ; Lipid-anchor ; Cytoplasmic side . Endoplasmic reticulum membrane . Golgi apparatus membrane . Late endosome. Cytoplasmic vesicle, phagosome membrane ; Lipid-anchor ; Cytoplasmic side . Cytoplasmic vesicle, phagosome. Colocalizes with OSBPL1A at the late endosome. Recruited to phagosomes containing S.aureus or M.tuberculosis.

Protein Name



Anti-Rab9 Picoband Antibody - Background

Ras-related protein Rab-9A, also called RAB9, is a protein that in humans is encoded by the RAB9A gene. This gene is mapped to Xp22.2. RAB9 has been localized to components of the endocytic/exocytic pathway and has been implicated in recycling of membrane receptors. It has been found that downregulation of RAB9A gene expression in HeLa cells induced severe cell vacuolation. RAB9A GTPase is directly bound by TIP47 in its active, GTP-bound conformation. Moreover, RAB9A increases the affinity of TIP47 for its cargo. What's more, this gene may involved in the transport of proteins between the endosomes and the trans Golgi network. RAB9A has been shown to interact with RABEPK and TIP47.

Ras-related protein Rab-9A

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human Rab9 (156-193aa KDATNVAAAFEEAVRRVLATEDRS DHLIQTDTVNLHRK), different from the related mouse sequence by three amino acids, and from the related rat sequence by two amino acids.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins.

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-Rab9 Picoband Antibody - Protein Information

Name RAB9A

Synonyms RAB9

Function

Involved in the transport of proteins between the endosomes and the trans Golgi network. Involved in the recruitment of SGSM2 to melanosomes and is required for the proper trafficking of melanogenic enzymes TYR, TYRP1 and DCT/TYRP2 to melanosomes in melanocytes.

Cellular Location

Cell membrane; Lipid-anchor; Cytoplasmic side. Endoplasmic reticulum membrane. Golgi apparatus membrane. Late endosome. Cytoplasmic vesicle, phagosome membrane; Lipid-anchor; Cytoplasmic side. Cytoplasmic vesicle, phagosome. Cytoplasmic vesicle membrane

{ECO:0000250|UniProtKB:Q9ROM6}.
Melanosome
{ECO:0000250|UniProtKB:Q9ROM6}.
Note=Colocalizes with OSBPL1A at the late
endosome (PubMed:16176980). Recruited
to phagosomes containing S.aureus or
M.tuberculosis (PubMed:21255211)

Anti-Rab9 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)