

Goat Anti-MOG Antibody

Peptide-affinity purified goat antibody
Catalog # AF1676a

Specification

Goat Anti-MOG Antibody - Product Information

| | |
|-------------------------|--|
| Application | WB |
| Primary Accession | Q16653 |
| Other Accession | NP_001008229 , 4340 |
| Reactivity Predicted | Human Mouse, Rat, Pig, Cow |
| Host | Goat |
| Clonality | Polyclonal |
| Concentration | 0.5 mg/ml |
| Isotype | IgG |
| Calculated MW | 28193 |

Goat Anti-MOG Antibody - Additional Information

Gene ID 4340

Other Names

Myelin-oligodendrocyte glycoprotein, MOG

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-MOG Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-MOG Antibody - Protein Information

Name MOG

Function

Mediates homophilic cell-cell adhesion (By



AF1676a staining (0.03 µg/ml) of Human Brain lysate (RIPA buffer, 30 µg total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.

Goat Anti-MOG Antibody - Background

The product of this gene is a membrane protein expressed on the oligodendrocyte cell surface and the outermost surface of myelin sheaths. Due to this localization, it is a primary target antigen involved in immune-mediated demyelination. This protein may be involved in completion and maintenance of the myelin sheath and in cell-cell communication. Alternatively spliced transcript variants encoding different isoforms have been identified.

Goat Anti-MOG Antibody - References

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. A major histocompatibility Class I locus contributes to multiple sclerosis susceptibility independently from

similarity). Minor component of the myelin sheath. May be involved in completion and/or maintenance of the myelin sheath and in cell-cell communication.

Cellular Location

[Isoform 1]: Cell membrane; Multi- pass membrane protein [Isoform 2]: Cell membrane; Single- pass type I membrane protein [Isoform 4]: Cell membrane; Single- pass type I membrane protein [Isoform 7]: Cell membrane; Single- pass type I membrane protein [Isoform 9]: Cell membrane; Single- pass type I membrane protein

Tissue Location

Found exclusively in the CNS, where it is localized on the surface of myelin and oligodendrocyte cytoplasmic membranes

HLA-DRB1*15:01. Cree BA, et al. PLoS One, 2010 Jun 25. PMID 20593013. Proteome analysis of the thalamus and cerebrospinal fluid reveals glycolysis dysfunction and potential biomarkers candidates for schizophrenia. Martins-de-Souza D, et al. J Psychiatr Res, 2010 May 14. PMID 20471030. The association of myelin oligodendrocyte glycoprotein gene and white matter volume in obsessive-compulsive disorder. Atmaca M, et al. J Affect Disord, 2010 Aug. PMID 20452030. Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614.

Goat Anti-MOG Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)