

ARID3B antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10626

Specification

ARID3B antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8IVW6
Other Accession	NM_006465 , NP_006456
Reactivity	Human, Mouse, Rat, Zebrafish, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Pig, Chicken, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	60kDa KDa

ARID3B antibody - C-terminal region - Additional Information

Gene ID 10620

Alias Symbol **BDP, DRIL2**

Other Names

AT-rich interactive domain-containing protein 3B, ARID domain-containing protein 3B, Bright and dead ringer protein, Bright-like protein, ARID3B, BDP, DRIL2

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ARID3B antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ARID3B antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ARID3B antibody - C-terminal region - Protein Information

Name ARID3B

Synonyms BDP, DRIL2

Function

Transcription factor which may be involved in neuroblastoma growth and malignant transformation. Favors nuclear targeting of ARID3A.

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00355, ECO:0000269|PubMed:10446990,

ECO:0000269|PubMed:17400556}

Tissue Location

Expressed in placenta, testis and leukocytes. Expressed in neuroblastoma. Present in K-562 erythrocytic leukemia cell line (at protein level).

ARID3B antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)