

**ZBTB40 antibody - N-terminal region**  
**Rabbit Polyclonal Antibody**  
Catalog # AI10630

**Specification**

**ZBTB40 antibody - N-terminal region - Product Information**

Application	WB
Primary Accession	<a href="#">Q9NUA8</a>
Other Accession	<a href="#">NM_014870</a> , <a href="#">NP_055685</a>
Reactivity	<b>Human, Rat, Pig</b>
Predicted	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>138kDa KDa</b>

**ZBTB40 antibody - N-terminal region - Additional Information**

**Gene ID** 9923

Alias Symbol	<b>KIAA0478,</b> <b>MGC133098,</b> <b>ZNF923</b>
--------------	--

**Other Names**

Zinc finger and BTB domain-containing protein 40, ZBTB40, KIAA0478

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**


Add 50 ul of distilled water. Final anti-ZBTB40 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

ZBTB40 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**ZBTB40 antibody - N-terminal region - Protein Information**

**Name** ZBTB40



**Synonyms** KIAA0478**Function**

May be involved in transcriptional regulation.

**Cellular Location**

Nucleus.

**ZBTB40 antibody - N-terminal region -  
Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)