

ZNF141 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10868**Specification****ZNF141 antibody - N-terminal region - Product Information**

| | |
|-------------------|--|
| Application | WB |
| Primary Accession | Q15928 |
| Other Accession | NM_003441 , NP_003432 |
| Reactivity | Human |
| Predicted | Human |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 55kDa KDa |

ZNF141 antibody - N-terminal region - Additional Information**Gene ID 7700**

| | |
|--|----------------------|
| Alias Symbol | D4S90, pHZ-44 |
| Other Names | |
| Zinc finger protein 141, ZNF141, D4S90 | |

Format


Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ZNF141 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ZNF141 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF141 antibody - N-terminal region - Protein Information**Name** ZNF141**Synonyms** D4S90**ZNF141 antibody - N-terminal region - References**

Varrault,A., (1998) Proc. Natl. Acad. Sci. U.S.A. 95 (15), 8835-8840 Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

Function

May be involved in transcriptional regulation as a repressor. Plays a role in limb development.

Cellular Location

Nucleus.

Tissue Location

Ubiquitously low expression.

**ZNF141 antibody - N-terminal region -
Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)