

AES antibody - C-terminal region

Rabbit Polyclonal Antibody Catalog # Al10997

Specification

AES antibody - C-terminal region - Product Information

Application WB
Primary Accession Other Accession NP 001121

Reactivity Human, Mouse,

Rat, Bovine

Predicted Human, Mouse,

Rat, Bovine, Dog

Host Rabbit
Clonality Polyclonal
Calculated MW 22kDa KDa

AES antibody - C-terminal region - Additional Information

Gene ID 166

Alias Symbol AES-1, AES-2,

ESP1, GRG, GRG5,

TLE5

Other Names

Amino-terminal enhancer of split, Amino enhancer of split, Gp130-associated protein GAM, Grg-5, Groucho-related protein 5, Protein ESP1, Protein GRG, AES, GRG, GRG5

Format


Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-AES antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

AES antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.



WB Suggested Anti-AES Antibody Titration:


 $2.5\mu g/ml$

Positive Control: Jurkat cell lysate

AES is supported by BioGPS gene expression data to be expressed in Jurkat

AES antibody - C-terminal region - References

Jan,Y., (2004) Cell 116 (5), 751-762Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.





AES antibody - C-terminal region - Protein Information

Name TLE5 (HGNC:307)

Synonyms AES, GRG, GRG5

Function

Transcriptional corepressor. Acts as dominant repressor towards other family members. Inhibits NF-kappa-B-regulated gene expression. May be required for the initiation and maintenance of the differentiated state. Essential for the transcriptional repressor activity of SIX3 during retina and lens development.

Cellular Location Nucleus.

Tissue Location

Found predominantly in muscle, heart and Placenta. In fetal tissues, abundantly expressed in the heart, lung, kidney, brain and liver

AES antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture