

**ZNF564 antibody - C-terminal region**  
**Rabbit Polyclonal Antibody**  
Catalog # AI11630

**Specification**

**ZNF564 antibody - C-terminal region - Product Information**

Application	WB
Primary Accession	<a href="#">Q8TBZ8</a>
Other Accession	<a href="#">NM_144976</a> , <a href="#">NP_659413</a>
Reactivity	Human, Mouse, Rat, Zebrafish, Pig, Horse, Bovine, Dog
Predicted	Mouse, Rat, Zebrafish, Pig, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	64kDa KDa

**ZNF564 antibody - C-terminal region - Additional Information**

**Gene ID** 163050

**Alias Symbol** **MGC26914**

**Other Names**

Zinc finger protein 564, ZNF564

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.


**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-ZNF564 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

ZNF564 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**ZNF564 antibody - C-terminal region - Protein Information**



**ZNF564 antibody - C-terminal region - References**

Isogai,T. Unpublished (2002)Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

**Name** ZNF564

**Function**

May be involved in transcriptional regulation.

**Cellular Location**

Nucleus.

**ZNF564 antibody - C-terminal region -  
Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)