

FBX22 Antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI16151**Specification****FBX22 Antibody - middle region - Product Information**

Application	WB
Primary Accession	O8NEZ5
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	44kDa KDa

FBX22 Antibody - middle region - Additional Information**Gene ID** 26263**Alias Symbol** **FBXO22, FBX22,**
Other NamesF-box only protein 22, F-box protein
FBX22p44, FBXO22, FBX22**Format**


Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & StorageAdd 50 μ l of distilled water. Final Anti-FBX22 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.**Precautions**

FBX22 Antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

FBX22 Antibody - middle region - Protein Information**Name** FBXO22**Synonyms** FBX22**Function**

Substrate-recognition component of the SCF



Host: Rabbit
Target Name: FBX22
Sample Tissue: NCI-H226 Whole Cell lysates
Antibody Dilution: 1.0µg/ml

FBX22 Antibody - middle region - Background

Substrate-recognition component of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex. Promotes the proteasome-dependent degradation of key sarcomeric proteins, such as alpha-actinin (ACTN2) and filamin-C (FLNC), essential for maintenance of normal contractile function.

FBX22 Antibody - middle region - ReferencesTan P., et al. Submitted (JUL-2000) to the EMBL/GenBank/DDBJ databases.
Ota T., et al. Nat. Genet. 36:40-45(2004).
Zody M.C., et al. Nature 440:671-675(2006).
Cenciarelli C., et al. Curr. Biol. 9:1177-1179(1999).
Gauci S., et al. Anal. Chem. 81:4493-4501(2009).

(SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex. Promotes the proteasome-dependent degradation of key sarcomeric proteins, such as alpha-actinin (ACTN2) and filamin-C (FLNC), essential for maintenance of normal contractile function.

Cellular Location

Cytoplasm, myofibril, sarcomere, Z line

Tissue Location

Predominantly expressed in liver, also enriched in cardiac muscle.

**FBX22 Antibody - middle region -
Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)