




**F2RL3 / PAR4 Antibody (Extracellular Domain)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS10331****Specification****F2RL3 / PAR4 Antibody (Extracellular Domain) -  
Product Information**

Application	<b>IHC</b>
Primary Accession	<a href="#">Q96R10</a>
Reactivity	<b>Human, Monkey</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>41kDa KDa</b>

**F2RL3 / PAR4 Antibody (Extracellular Domain) -  
Additional Information****Gene ID 9002****Other Names**Proteinase-activated receptor 4, PAR-4,  
Coagulation factor II receptor-like 3,  
Thrombin receptor-like 3, F2RL3, PAR4**Target/Specificity**Human F2RL3 / PAR4. BLAST analysis of the  
peptide immunogen showed no homology  
with other human proteins.**Reconstitution & Storage**

Long term: -70°C; Short term: +4°C

**Precautions**F2RL3 / PAR4 Antibody (Extracellular  
Domain) is for research use only and not for  
use in diagnostic or therapeutic procedures.**F2RL3 / PAR4 Antibody (Extracellular Domain) -  
Protein Information****Name F2RL3****Synonyms PAR4****Function**Receptor for activated thrombin or trypsin  
coupled to G proteins that stimulate  
phosphoinositide hydrolysis. May play a role  
in platelets activation.Anti-F2RL3 / PAR4 antibody ALS10331 IHC of  
human bone marrow.Anti-F2RL3 / PAR4 antibody IHC of human  
Ovary, Carcinoma.

**Cellular Location**

Cell membrane; Multi-pass membrane protein.

**Tissue Location**

Widely expressed, with highest levels in lung, pancreas, thyroid, testis and small intestine. Not expressed in brain, kidney, spinal cord and peripheral blood leukocytes. Also detected in platelets

**Volume**

50 µl

**F2RL3 / PAR4 Antibody (Extracellular Domain) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-F2RL3 / PAR4 antibody IHC of human Lymph Node, Hodgkins Lymphoma.

**F2RL3 / PAR4 Antibody (Extracellular Domain) - Background**

Receptor for activated thrombin or trypsin coupled to G proteins that stimulate phosphoinositide hydrolysis. May play a role in platelets activation.

**F2RL3 / PAR4 Antibody (Extracellular Domain) - References**

Kahn M.L.,et al.J. Biol. Chem. 273:23290-23296(1998).  
Kahn M.L.,et al.Nature 394:690-694(1998).  
Xu W.-F.,et al.Proc. Natl. Acad. Sci. U.S.A. 95:6642-6646(1998).  
King M.M.,et al.Submitted (OCT-2003) to the EMBL/GenBank/DDBJ databases.  
Kahn M.L.,et al.J. Clin. Invest. 103:879-887(1999).