

TIRAP Antibody (C-Terminus)
Rabbit Polyclonal Antibody
Catalog # ALS11722**Specification****TIRAP Antibody (C-Terminus) - Product Information**

Application	IHC
Primary Accession	P58753
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	24kDa KDa

TIRAP Antibody (C-Terminus) - Additional Information**Gene ID** 114609**Other Names**

Toll/interleukin-1 receptor domain-containing adapter protein, TIR domain-containing adapter protein, Adaptor protein Wyatt, MyD88 adapter-like protein, MyD88-2, TIRAP, MAL

Target/Specificity

peptide corresponding to 15 amino acids near the C-terminus of mouse TIRAP

Reconstitution & Storage


Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.

Precautions


TIRAP Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

TIRAP Antibody (C-Terminus) - Protein Information**Name** TIRAP**Synonyms** MAL**Function**

Adapter involved in TLR2 and TLR4



Anti-TIRAP antibody IHC of human heart.



Anti-TIRAP antibody IHC of human kidney.

TIRAP Antibody (C-Terminus) - Background

Adapter involved in TLR2 and TLR4 signaling pathways in the innate immune response. Acts via IRAK2 and TRAF-6, leading to the activation of NF-kappa-B, MAPK1, MAPK3 and JNK, and resulting in cytokine secretion and the inflammatory response. Positively regulates the production of TNF-alpha and interleukin-6.

TIRAP Antibody (C-Terminus) - References

Fitzgerald K.A., et al. Nature 413:78-83(2001).

signaling pathways in the innate immune response. Acts via IRAK2 and TRAF-6, leading to the activation of NF-kappa-B, MAPK1, MAPK3 and JNK, and resulting in cytokine secretion and the inflammatory response. Positively regulates the production of TNF-alpha and interleukin-6.

Cellular Location

Cytoplasm. Cell membrane. Membrane.
Note=Colocalizes with DAB2IP at the plasma membrane

Tissue Location

Highly expressed in liver, kidney, spleen, skeletal muscle and heart. Also detected in peripheral blood leukocytes, lung, placenta, small intestine, thymus, colon and brain

Hornig T.,et al.Nat. Immunol. 2:835-841(2001).
Kirk P.B.,et al.Submitted (AUG-2001) to the EMBL/GenBank/DDBJ databases.
Hardy M.P.,et al.Submitted (MAR-2004) to the EMBL/GenBank/DDBJ databases.
Nakajima T.,et al.Immunogenetics 60:727-735(2008).

TIRAP Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)