

RTN4 / Nogo Antibody (Internal)
Goat Polyclonal Antibody
Catalog # ALS13214

Specification

RTN4 / Nogo Antibody (Internal) - Product Information

Application	IHC
Primary Accession	Q9NOC3
Reactivity	Human, Monkey
Host	Goat
Clonality	Polyclonal
Calculated MW	130kDa KDa

RTN4 / Nogo Antibody (Internal) - Additional Information

Gene ID 57142

Other Names

Reticulon-4, Foccen, Neurite outgrowth inhibitor, Nogo protein, Neuroendocrine-specific protein, NSP, Neuroendocrine-specific protein C homolog, RTN-x, Reticulon-5, RTN4, KIAA0886, NOGO

Target/Specificity

Human RTN4 / Nogo. This antibody is expected to recognize reported isoform A (NP_065393.1) and isoform E (NP_997404.1).

Reconstitution & Storage

Store at -20°C. Minimize freezing and thawing.

Precautions


RTN4 / Nogo Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

RTN4 / Nogo Antibody (Internal) - Protein Information


Name RTN4 ([HGNC:14085](#))

Function

Required to induce the formation and stabilization of endoplasmic reticulum (ER) tubules (PubMed: <a href="http://www.unipr



Anti-RTN4 / Nogo antibody IHC of human brain, cortex.



Anti-RTN4 / Nogo antibody IHC of human testis.

RTN4 / Nogo Antibody (Internal) - Background

Developmental neurite growth regulatory factor with a role as a negative regulator of axon-axon adhesion and growth, and as a facilitator of neurite branching. Regulates neurite fasciculation, branching and extension in the developing nervous system. Involved in down-regulation of growth, stabilization of wiring and restriction of plasticity in the adult CNS. Regulates the radial migration of cortical

ot.org/citations/27619977" target="_blank">27619977, PubMed:25612671, PubMed:24262037). They regulate membrane morphogenesis in the ER by promoting tubular ER production (PubMed:27619977, PubMed:25612671, PubMed:24262037, PubMed:27786289). They influence nuclear envelope expansion, nuclear pore complex formation and proper localization of inner nuclear membrane proteins (PubMed:26906412). However each isoform have specific functions mainly depending on their tissue expression specificities (Probable).

Cellular Location

[Isoform A]: Endoplasmic reticulum membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein; Cytoplasmic side Note=Anchored to the membrane of the endoplasmic reticulum (ER) through 2 putative transmembrane domains. Localizes throughout the ER tubular network (PubMed:27619977). Co-localizes with TMEM33 at the ER sheets [Isoform C]: Endoplasmic reticulum membrane; Multi-pass membrane protein

Tissue Location

Isoform A: is specifically expressed in brain and testis and weakly in heart and skeletal muscle. Isoform B: widely expressed except for the liver. Highly expressed in endothelial cells and vascular smooth muscle cells, including blood vessels and mesenteric arteries (PubMed:15034570, PubMed:21183689). Isoform C: is expressed in brain, skeletal muscle and adipocytes. Isoform D is testis-specific.

neurons via an RTN4R-LINGO1 containing receptor complex (By similarity). Isoform 2 reduces the anti-apoptotic activity of Bcl-xl and Bcl-2. This is likely consecutive to their change in subcellular location, from the mitochondria to the endoplasmic reticulum, after binding and sequestration. Isoform 2 and isoform 3 inhibit BACE1 activity and amyloid precursor protein processing.

RTN4 / Nogo Antibody (Internal) - References

Yang J.,et al.Cytogenet. Cell Genet. 88:101-102(2000).
Prinjha R.,et al.Nature 403:383-384(2000).
Tagami S.,et al.Oncogene 19:5736-5746(2000).
Zhou Z.M.,et al.Reproduction 123:227-234(2002).
Oertle T.,et al.J. Mol. Biol. 325:299-323(2003).

**RTN4 / Nogo Antibody (Internal) -
Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)