

FURIN Antibody (aa740-790)
Rabbit Polyclonal Antibody
Catalog # ALS14903

Specification

FURIN Antibody (aa740-790) - Product Information

Application	IHC
Primary Accession	P09958
Reactivity	Human, Mouse, Rat, Monkey, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	87kDa KDa

FURIN Antibody (aa740-790) - Additional Information

Gene ID 5045

Other Names

Furin, 3.4.21.75, Dibasic-processing enzyme, Paired basic amino acid residue-cleaving enzyme, PACE, FURIN, FUR, PACE, PCSK3

Target/Specificity

Human FURIN

Reconstitution & Storage

Store at 4°C for short term applications. For long term storage, aliquot and store at -20°C.

Precautions

FURIN Antibody (aa740-790) is for research use only and not for use in diagnostic or therapeutic procedures.


FURIN Antibody (aa740-790) - Protein Information

Name FURIN

{ECO:0000303|PubMed:7690548, ECO:0000312|HGNC:HGNC:8568}

Function

Ubiquitous endoprotease within constitutive



Anti-FURIN antibody IHC of human adrenal.

FURIN Antibody (aa740-790) - Background

Furin is likely to represent the ubiquitous endoprotease activity within constitutive secretory pathways and capable of cleavage at the RX(K/R)R consensus motif.

FURIN Antibody (aa740-790) - References

- van den Ouweland A.M.W., et al. Nucleic Acids Res. 18:664-664(1990).
- Wise R.J., et al. Proc. Natl. Acad. Sci. U.S.A. 87:9378-9382(1990).
- Barr P.J., et al. DNA Cell Biol. 10:319-328(1991).
- Van den Ouweland A.M.W., et al. Nucleic Acids Res. 17:7101-7102(1989).
- Roebroek A.J.M., et al. EMBO J. 5:2197-2202(1986).

secretory pathways capable of cleavage at the RX(K/R)R consensus motif (PubMed:11799113, PubMed:1629222, PubMed:1713771, PubMed:2251280, PubMed:24666235, PubMed:25974265, PubMed:7592877, PubMed:7690548, PubMed:9130696). Mediates processing of TGFB1, an essential step in TGF-beta-1 activation (PubMed:7737999). Converts through proteolytic cleavage the non-functional Brain natriuretic factor prohormone into its active hormone BNP(1-32) (PubMed:20489134, PubMed:21763278).

Cellular Location

Golgi apparatus, trans-Golgi network membrane; Single-pass type I membrane protein. Cell membrane; Single-pass type I membrane protein. Secreted. Endosome membrane; Single-pass type I membrane protein. Note=Shuttles between the trans-Golgi network and the cell surface (PubMed:9412467, PubMed:11799113). Propeptide cleavage is a prerequisite for exit of furin molecules out of the endoplasmic reticulum (ER). A second cleavage within the propeptide occurs in the trans Golgi network (TGN), followed by the release of the propeptide and the

activation of furin (PubMed:11799113)

Tissue Location

Seems to be expressed ubiquitously.

FURIN Antibody (aa740-790) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)