

## DNAJC5 / CSP Antibody (aa177-191)

Goat Polyclonal Antibody Catalog # ALS15870

### **Specification**

DNAJC5 / CSP Antibody (aa177-191) - Product Information

Application WB, IHC Primary Accession Q9H3Z4

Reactivity Human, Mouse,

Rat, Rabbit,

Monkey, Chicken, Horse, Bovine, Guinea Pig, Dog

Host Goat
Clonality Polyclonal
Calculated MW 22kDa KDa

DNAJC5 / CSP Antibody (aa177-191) - Additional Information

Gene ID 80331

### **Other Names**

DnaJ homolog subfamily C member 5, Cysteine string protein, CSP, DNAJC5, CSP

Target/Specificity
Human DNAJC5 / CSP

**Reconstitution & Storage** Store at -20°C. Minimize freezing and thawing.

## **Precautions**


DNAJC5 / CSP Antibody (aa177-191) is for research use only and not for use in diagnostic or therapeutic procedures.

DNAJC5 / CSP Antibody (aa177-191) - Protein Information


Name DNAJC5 (HGNC:16235)

### Function

Acts as a general chaperone in regulated exocytosis (By similarity). Acts as a co-chaperone for the SNARE protein SNAP-25 (By similarity). Involved in the calcium-mediated control of a late stage of



DNAJC5 antibody (0.5 ug/ml) staining of Human Cerebellum lysate (35 ug protein in RIPA buffer).




Anti-DNAJC5 / CSP antibody IHC staining of human skeletal muscle.

# DNAJC5 / CSP Antibody (aa177-191) - Background

May have an important role in presynaptic function. May be involved in calcium-dependent neurotransmitter release at nerve endings (By similarity).

## DNAJC5 / CSP Antibody (aa177-191) - References





Tel: 858.875.1900 Fax: 858.622.0609

exocytosis (By similarity). May have an important role in presynaptic function. May be involved in calcium-dependent neurotransmitter release at nerve endings (By similarity).

### **Cellular Location**

Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q29455}. Membrane {ECO:0000250|UniProtKB:Q29455}; Lipid-anchor {ECO:0000250|UniProtKB:Q29455}. Cytoplasmic vesicle, secretory vesicle, chromaffin granule membrane {ECO:0000250|UniProtKB:Q29455}. Melanosome. Cell membrane. Note=The association with membranes is regulated by palmitoylation (By similarity). Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065). {ECO:0000250|UniProtKB:Q29455, ECO:0000269|PubMed:17081065}

### **Tissue Location**

Expressed in pancreas, kidney, skeletal muscle, liver, lung, placenta, brain and heart.

## DNAJC5 / CSP Antibody (aa177-191) -**Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Coppola T., et al. FEBS Lett. 391:269-272(1996). Ota T., et al. Nat. Genet. 36:40-45(2004). Deloukas P., et al. Nature 414:865-871(2001). Mural R.J., et al. Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases. Hattori A., et al. DNA Res. 7:357-366(2000).