



ENPP2 / Autotaxin Antibody (aa698-712)
Goat Polyclonal Antibody
Catalog # ALS16150**Specification****ENPP2 / Autotaxin Antibody (aa698-712) -
Product Information**

Application	WB, IHC
Primary Accession	O13822
Reactivity	Human, Monkey
Host	Goat
Clonality	Polyclonal
Calculated MW	99kDa KDa

**ENPP2 / Autotaxin Antibody (aa698-712) -
Additional Information****Gene ID 5168****Other Names**Ectonucleotide
pyrophosphatase/phosphodiesterase family
member 2, E-NPP 2, 3.1.4.39, Autotaxin,
Extracellular lysophospholipase D, LysoPLD,
ENPP2, ATX, PDNP2**Target/Specificity**Human ENPP2 / Autotaxin. This antibody is
expected to recognize all reported isoforms
(NP_006200.3; NP_001035181.1,
NP_001124335.1).**Reconstitution & Storage**Store at -20°C. Minimize freezing and
thawing.**Precautions**ENPP2 / Autotaxin Antibody (aa698-712) is
for research use only and not for use in
diagnostic or therapeutic procedures.**ENPP2 / Autotaxin Antibody (aa698-712) - Protein
Information****Name ENPP2****Function**Hydrolyzes lysophospholipids to produce
the signaling molecule lysophosphatidic
acid (LPA) in extracellular fluidsENPP2 antibody (0.3 ug/ml) staining of
Human Placenta lysate (35 ug protein in RIPA
buffer).Anti-ENPP2 / Autotaxin antibody IHC staining
of human brain, cortex.

(PubMed:15769751, PubMed:26371182, PubMed:27754931, PubMed:14500380, PubMed:12354767), Major substrate is lysophosphatidylcholine (PubMed:12176993, PubMed:27754931, PubMed:14500380). Also can act on sphingosylphosphorylcholine producing sphingosine-1-phosphate, a modulator of cell motility (PubMed:14500380). Can hydrolyze, in vitro, bis-pNPP, to some extent pNP-TMP, and barely ATP (PubMed:15769751, PubMed:12176993). Involved in several motility-related processes such as angiogenesis and neurite outgrowth. Acts as an angiogenic factor by stimulating migration of smooth muscle cells and microtubule formation (PubMed:11559573). Stimulates migration of melanoma cells, probably via a pertussis toxin-sensitive G protein (PubMed:1733949). May have a role in induction of parturition (PubMed:12176993). Possible involvement in cell proliferation and adipose tissue development (Probable). Tumor cell motility-stimulating factor (PubMed:15769751).

Anti-ENPP2 / Autotaxin antibody IHC staining of human liver.

ENPP2 / Autotaxin Antibody (aa698-712) - Background

Hydrolyzes lysophospholipids to produce lysophosphatidic acid (LPA) in extracellular fluids. Major substrate is lysophosphatidylcholine. Also can act on sphingosylphosphorylcholine producing sphingosine-1-phosphate, a modulator of cell motility. Can hydrolyze, in vitro, bis-pNPP, to some extent pNP-TMP, and barely ATP. Involved in several motility-related processes such as angiogenesis and neurite outgrowth. Acts as an angiogenic factor by stimulating migration of smooth muscle cells and microtubule formation. Stimulates migration of melanoma cells, probably via a pertussis toxin-sensitive G protein. May have a role in induction of parturition. Possible involvement in cell proliferation and adipose tissue development. Tumor cell motility-stimulating factor.

ENPP2 / Autotaxin Antibody (aa698-712) - References

- Murata J., et al. J. Biol. Chem. 269:30479-30484(1994).
- Kawagoe H., et al. Genomics 30:380-384(1995).
- Lee H.Y., et al. Biochem. Biophys. Res. Commun. 218:714-719(1996).
- van Meeteren L.A., et al. J. Biol. Chem. 280:21155-21161(2005).
- Giganti A., et al. J. Biol. Chem. 283:7776-7789(2008).

itations/1733949" target="_blank">1733949, PubMed:11559573). Required for LPA production in activated platelets, cleaves the sn-1 lysophospholipids to generate sn-1 lysophosphatidic acids containing predominantly 18:2 and 20:4 fatty acids (PubMed:21393252). Shows a preference for the sn-1 to the sn-2 isomer of 1-O-alkyl-sn-glycero-3-phosphocholine (lyso-PAF) (PubMed:21393252).

Cellular Location

Secreted

Tissue Location

Detected in blood plasma (at protein level) (PubMed:12176993, PubMed:26371182). Predominantly expressed in brain, placenta, ovary, and small intestine. Expressed in a number of carcinomas such as hepatocellular and prostate carcinoma, neuroblastoma and non-small-cell lung cancer. Expressed in body fluids such as plasma, cerebral spinal fluid (CSF), saliva, follicular and amniotic fluids. Not detected in leukocytes. Isoform 1 is more highly expressed in peripheral tissues than in the central nervous system (CNS) Adipocytes only express isoform 1. Isoform 3 is more highly expressed in the brain than in peripheral tissues.

ENPP2 / Autotaxin Antibody (aa698-712) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)