

Ki67 Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1253a

Specification

Ki67 Antibody - Product Information

Application	WB
Primary Accession	P46013
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1

Description

Ki67: antigen identified by monoclonal antibody Ki-67. Ki67 antigen is the prototypic cell cycle related nuclear protein, expressed by proliferating cells in all phases of the active cell cycle (G1, S, G2 and M phase). It is absent in resting (G0) cells. Ki67 antibodies are useful in establishing the cell growing fraction in neoplasms (immunohistochemically quantified by determining the number of Ki67 positive cells among the total number of resting cells = Ki67 index). In neoplastic tissues the prognostic value is comparable to the tritiated thymidine labelling index. The correlation between low Ki67 index and histologically low grade tumours is strong. Ki67 is routinely used as a neuronal marker of cell cycling and proliferation.

Immunogen

Purified recombinant fragment of Ki167 (aa3118-3256) expressed in E. Coli.

Formulation

Ascitic fluid containing 0.03% sodium azide.

Ki67 Antibody - Additional Information

Gene ID 4288

Other Names

Antigen KI-67, MKI67

Dilution

WB ~ 1/500 - 1/2000

Storage




Figure 1: Western blot analysis using Ki67 mouse mAb against truncated Trx-Ki67 recombinant protein(1), truncated Ki67 (aa3118-3256)-His recombinant protein(2) and truncated Ki67 (aa3118-3256)-hlgGfc transfected CHO-K1 cell lysate(3).

Ki67 Antibody - References

1. Folia Histochem Cytobiol. 2007;45(4):357-66.
2. Tumori. 2008 May-Jun;94(3):389-97.

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Ki67 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Ki67 Antibody - Protein Information

Name MKI67 ([HGNC:7107](#))

Function

Required to maintain individual mitotic chromosomes dispersed in the cytoplasm following nuclear envelope disassembly (PubMed:27362226). Associates with the surface of the mitotic chromosome, the perichromosomal layer, and covers a substantial fraction of the chromosome surface (PubMed:27362226). Prevents chromosomes from collapsing into a single chromatin mass by forming a steric and electrostatic charge barrier: the protein has a high net electrical charge and acts as a surfactant, dispersing chromosomes and enabling independent chromosome motility (PubMed:27362226). Binds DNA, with a preference for supercoiled DNA and AT-rich DNA (PubMed:10878551). Does not contribute to the internal structure of mitotic chromosomes (By similarity). May play a role in chromatin organization (PubMed:24867636). It is however unclear whether it plays a direct role in chromatin organization or whether it is an indirect consequence of its function in maintaining mitotic chromosomes dispersed (Probable).

Cellular Location

Chromosome. Nucleus. Nucleus, nucleolus
Note=Associates with the surface of the mitotic chromosome, the perichromosomal

layer, and covers a substantial fraction of the mitotic chromosome surface (PubMed:27362226). Associates with satellite DNA in G1 phase (PubMed:9510506). Binds tightly to chromatin in interphase, chromatin-binding decreases in mitosis when it associates with the surface of the condensed chromosomes (PubMed:15896774, PubMed:22002106). Predominantly localized in the G1 phase in the perinucleolar region, in the later phases it is also detected throughout the nuclear interior, being predominantly localized in the nuclear matrix (PubMed:22002106).

Ki67 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)