

**NT5E Antibody**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO1306a**

**Specification**

**NT5E Antibody - Product Information**

Application	<b>IHC, WB</b>
Primary Accession	<a href="#">P21589</a>
Reactivity	<b>Human</b>
Host	<b>Mouse</b>
Clonality	<b>Monoclonal</b>
Isotype	<b>IgG1</b>
Calculated MW	<b>70kDa KDa</b>

**Description**

5'-nucleotidase, ecto (NT5E), also known as CD73 (Cluster of Differentiation 73). Ecto-5-prime-nucleotidase (5-prime-ribonucleotide phosphohydrolase; EC 3.1.3.5) catalyzes the conversion at neutral pH of purine 5-prime mononucleotides to nucleosides, the preferred substrate being AMP. The enzyme consists of a dimer of 2 identical 70-kD subunits bound by a glycosyl phosphatidyl inositol linkage to the external face of the plasma membrane. The enzyme is used as a marker of lymphocyte differentiation. Consequently, a deficiency of NT5 occurs in a variety of immunodeficiency diseases (e.g., see MIM 102700, MIM 300300). Other forms of 5-prime nucleotidase exist in the cytoplasm and lysosomes and can be distinguished from ecto-NT5 by their substrate affinities, requirement for divalent magnesium ion, activation by ATP, and inhibition by inorganic phosphate.

**Immunogen**

Purified recombinant fragment of NT5E expressed in E. Coli.

**Formulation**

Ascitic fluid containing 0.03% sodium azide.

**NT5E Antibody - Additional Information**

**Gene ID 4907**

**Other Names**

5'-nucleotidase, 5'-NT, 3.1.3.5,




Figure 1: Immunohistochemical analysis of paraffin-embedded human lung cancer (A), cholangiocarcinoma (B), lymph node (C) and esophagus (D) tissues using NT5E mouse mAb with DAB staining.




Figure 1: Western blot analysis using FRK mouse mAb against K562 cell lysate (1).

**NT5E Antibody - References**

1. Oncol Rep. 2007 Jun;17(6):1341-6.
2. Neurochem Int. 2003 Dec;43(7):621-8.

Ecto-5'-nucleotidase, CD73, NT5E, NT5, NTE

**Dilution**

IHC~~1/200 - 1/1000

WB~~1:500~~2000

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

NT5E Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**NT5E Antibody - Protein Information**

**Name** NT5E

**Synonyms** NT5, NTE

**Function**

Hydrolyzes extracellular nucleotides into membrane permeable nucleosides. Exhibits AMP-, NAD-, and NMN-nucleosidase activities.

**Cellular Location**

Cell membrane; Lipid-anchor, GPI-anchor.

**NT5E Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)