

Mouse Monoclonal Antibody to TLR9
Purified Mouse Monoclonal Antibody
Catalog # A02370a

Specification

Mouse Monoclonal Antibody to TLR9 - Product Information

Application	E, WB, FC, ICC, IHC
Primary Accession	Q9NR96
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse IgG2a
Calculated MW	115.8kDa KDa

Description

The protein encoded by this gene is a member of the Toll-like receptor (TLR) family which plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from Drosophila to humans and share structural and functional similarities. They recognize pathogen-associated molecular patterns (PAMPs) that are expressed on infectious agents, and mediate the production of cytokines necessary for the development of effective immunity. The various TLRs exhibit different patterns of expression. This gene is preferentially expressed in immune cell rich tissues, such as spleen, lymph node, bone marrow and peripheral blood leukocytes. Studies in mice and human indicate that this receptor mediates cellular response to unmethylated CpG dinucleotides in bacterial DNA to mount an innate immune response.;

Immunogen


Purified recombinant fragment of human TLR9 (AA: 868-1016) expressed in E. Coli.

Formulation


Purified antibody in PBS with 0.05% sodium azide

Application Note

ELISA: 1/10000; WB: 1/500 - 1/2000; IHC: 1/200 - 1/1000; ICC: 1/200 - 1/1000; FCM: 1/200 - 1/400



Black line: Control Antigen (100 ng); Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)



Western blot analysis using TLR9 mAb against human TLR9 (AA: 868-1016) recombinant protein. (Expected MW is 43.4 kDa)

Mouse Monoclonal Antibody to TLR9 - Additional Information

Gene ID 54106

Other Names
CD289

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mouse Monoclonal Antibody to TLR9 is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Monoclonal Antibody to TLR9 - Protein Information


Name TLR9

Function


Key component of innate and adaptive immunity. TLRs (Toll-like receptors) control host immune response against pathogens through recognition of molecular patterns specific to microorganisms. TLR9 is a nucleotide-sensing TLR which is activated by unmethylated cytidine-phosphate-guanosine (CpG) dinucleotides. Acts via MYD88 and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response (PubMed:11564765, PubMed:17932028). Controls lymphocyte response to Helicobacter infection (By similarity). Upon CpG stimulation, induces B-cell proliferation, activation, survival and antibody production (PubMed:23857366).

Cellular Location


Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q9EQU3};
Single-pass type I membrane protein {ECO:0000250|UniProtKB:Q9EQU3}.
Endosome {ECO:0000250|UniProtKB:Q9EQU3}.



Western blot analysis using TLR9 mAb against HEK293 (1) and TLR9 (AA: 868-1016)-hlgGfc transfected HEK293 (2) cell lysate.



Flow cytometric analysis of A549 cells using TLR9 mouse mAb (green) and negative control (red).



Lysosome

{ECO:0000250|UniProtKB:Q9EQU3}.
Cytoplasmic vesicle, phagosome
{ECO:0000250|UniProtKB:Q9EQU3}.
Note=Relocalizes from endoplasmic reticulum to endosome and lysosome upon stimulation with agonist. Exit from the ER requires UNC93B1. Endolysosomal localization is required for proteolytic cleavage and subsequent activation. Intracellular localization of the active receptor may prevent from responding to self nucleic acid.
{ECO:0000250|UniProtKB:Q9EQU3}

Tissue Location


Highly expressed in spleen, lymph node, tonsil and peripheral blood leukocytes, especially in plasmacytoid pre-dendritic cells. Levels are much lower in monocytes and CD11c+ immature dendritic cells. Also detected in lung and liver

Mouse Monoclonal Antibody to TLR9 - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Immunofluorescence analysis of SK-OV-3 cells using TLR9 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin. Secondary antibody from Fisher



Immunohistochemical analysis of paraffin-embedded ovarian cancer tissues using TLR9 mouse mAb with DAB staining.



Immunohistochemical analysis of paraffin-embedded endometrial cancer tissues using TLR9 mouse mAb with DAB staining.

Mouse Monoclonal Antibody to TLR9 - References

1. J Virol. 2015 Nov;89(22):11396-405. ;
2. Immunogenetics. 2014 Dec;66(12):675-81.;