

**ABCC3 Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP10144C**

**Specification**

**ABCC3 Antibody (Center) - Product Information**

Application **WB, IHC-P, FC,E**  
Primary Accession [O15438](#)  
Other Accession [NP\\_003777.2](#),  
[NP\\_001137542.1](#)  
Reactivity **Human**  
Host **Rabbit**  
Clonality **Polyclonal**  
Isotype **Rabbit Ig**  
Antigen Region **899-925**

**ABCC3 Antibody (Center) - Additional Information**

**Gene ID 8714**

**Other Names**

Canalicular multispecific organic anion transporter 2, ATP-binding cassette sub-family C member 3, Multi-specific organic anion transporter D, MOAT-D, Multidrug resistance-associated protein 3, ABCC3, CMOAT2, MLP2, MRP3

**Target/Specificity**

This ABCC3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 899-925 amino acids from the Central region of human ABCC3.

**Dilution**


WB~~1:1000  
IHC-P~~1:10~50  
FC~~1:10~50

**Format**


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw



All lanes : Anti-ABCC3 Antibody (Center) at 1:1000 dilution Lane 1: Human brain lysate Lane 2: Human testis lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 169 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-ABCC3 Antibody (Center) at 1:1000 dilution Lane 1: HepG2 whole cell lysate Lane 2: human liver lysate

cycles.

### Precautions

ABCC3 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

### ABCC3 Antibody (Center) - Protein Information


Name ABCC3 ([HGNC:54](#))

Synonyms CMOAT2, MLP2, MRP3


### Function

ATP-dependent transporter of the ATP-binding cassette (ABC) family that bind and hydrolyze ATP to enable active transport of various substrates including many drugs, toxicants and endogenous compound across cell membranes (PubMed:[11581266](http://www.uniprot.org/citations/11581266) target="\_blank">11581266</a>, PubMed:[15083066](http://www.uniprot.org/citations/15083066) target="\_blank">15083066</a>, PubMed:[10359813](http://www.uniprot.org/citations/10359813) target="\_blank">10359813</a>). Transports glucuronide conjugates such as bilirubin diglucuronide, estradiol-17-beta-o-glucuronide and GSH conjugates such as leukotriene C4 (LTC4) (PubMed:[15083066](http://www.uniprot.org/citations/15083066) target="\_blank">15083066</a>, PubMed:[11581266](http://www.uniprot.org/citations/11581266) target="\_blank">11581266</a>). Transports also various bile salts (taurocholate, glycocholate, taurochenodeoxycholate-3-sulfate, tauroolithocholate- 3-sulfate) (By similarity). Does not contribute substantially to bile salt physiology but provides an alternative route for the export of bile acids and glucuronides from cholestatic hepatocytes (By similarity). Can confers resistance to various anticancer drugs, methotrexate, tenoposide and etoposide, by decreasing accumulation of these drugs in cells (PubMed:[11581266](http://www.uniprot.org/citations/11581266) target="\_blank">11581266</a>, PubMed:[10359813](http://www.uniprot.org/citations/10359813) target="\_blank">10359813</a>).

Lysates/proteins at 20 µg per lane.  
Secondary Goat Anti-Rabbit IgG, (H+L),  
Peroxidase conjugated at 1/10000 dilution.  
Predicted band size : 169 kDa  
Blocking/Dilution buffer: 5% NFDN/TBST.



ABCC3 antibody (Center) (Cat. #AP10144c) immunohistochemistry analysis in formalin fixed and paraffin embedded human colon carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the ABCC3 antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



ABCC3 Antibody (Center) (Cat. #AP10144c) flow cytometric analysis of MDA-MB435 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

### ABCC3 Antibody (Center) - Background

**Cellular Location**

Basolateral cell membrane; Multi-pass membrane protein

**Tissue Location**

Mainly expressed in the liver. Also expressed in small intestine, colon, prostate, testis, brain and at a lower level in the kidney.

**ABCC3 Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the MRP subfamily which is involved in multi-drug resistance. The specific function of this protein has not yet been determined; however, this protein may play a role in the transport of biliary and intestinal excretion of organic anions. Alternatively spliced variants which encode different protein isoforms have been described; however, not all variants have been fully characterized.

**ABCC3 Antibody (Center) - References**

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)  
Hoffman, A.D., et al. Protein J. 29(5):373-379(2010)  
Liu, C.Y., et al. Carcinogenesis 31(7):1259-1263(2010)  
Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :  
Moyer, A.M., et al. Cancer Epidemiol. Biomarkers Prev. 19(3):811-821(2010)