

**GPAA1 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP10722a**

**Specification**

**GPAA1 Antibody (N-term) - Product Information**

Application	<b>WB, IHC-P, FC,E</b>
Primary Accession	<a href="#">O43292</a>
Other Accession	<a href="#">NP_003792.1</a>
Reactivity	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>Rabbit Ig</b>
Calculated MW	<b>67623</b>
Antigen Region	<b>46-73</b>

**GPAA1 Antibody (N-term) - Additional Information**

**Gene ID** 8733

**Other Names**

Glycosylphosphatidylinositol anchor attachment 1 protein, GPI anchor attachment protein 1, GAA1 protein homolog, hGAA1, GPAA1, GAA1

**Target/Specificity**

This GPAA1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 46-73 amino acids from the N-terminal region of human GPAA1.

**Dilution**


WB~~1:1000  
IHC-P~~1:50~100  
FC~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.



GPAA1 Antibody (N-term) (Cat. #AP10722a) western blot analysis in U251 cell line lysates (35ug/lane). This demonstrates the GPAA1 antibody detected the GPAA1 protein (arrow).



GPAA1 antibody (N-term) (Cat. #AP10722a) immunohistochemistry analysis in formalin fixed and paraffin embedded human skeletal muscle followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the GPAA1 antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

### Precautions

GPAA1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

### GPAA1 Antibody (N-term) - Protein Information

**Name** GPAA1

**Synonyms** GAA1

### Function

Essential for GPI-anchoring of precursor proteins but not for GPI synthesis. Acts before or during formation of the carbonyl intermediate.

### Cellular Location

Endoplasmic reticulum membrane;  
Multi-pass membrane protein


### Tissue Location

Ubiquitously expressed in fetal and adult tissues. Expressed at higher levels in fetal tissues than adult tissues

### GPAA1 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



### GPAA1 Antibody (N-term) - Background

Posttranslational glycosylphosphatidylinositol (GPI) anchor attachment serves as a general mechanism for linking proteins to the cell surface membrane. The protein encoded by this gene presumably functions in GPI anchoring at the GPI transfer step. The mRNA transcript is ubiquitously expressed in both fetal and adult tissues. The anchor attachment protein 1 contains an N-terminal signal sequence, 1 cAMP- and cGMP-dependent protein kinase phosphorylation site, 1 leucine zipper pattern, 2 potential N-glycosylation sites, and 8 putative transmembrane domains.

### GPAA1 Antibody (N-term) - References

Jiang, W.W., et al. Mol. Cancer 6, 74 (2007) :  
Olsen, J.V., et al. Cell 127(3):635-648(2006)  
Ho, J.C., et al. Int. J. Cancer 119(6):1330-1337(2006)  
Vainauskas, S., et al. J. Biol. Chem. 280(16):16402-16409(2005)  
Vainauskas, S., et al. J. Biol. Chem. 279(8):6540-6545(2004)