

**COL8A2 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP11859B**

**Specification**

**COL8A2 Antibody (C-term) - Product Information**

Application	<b>IF, WB, IHC-P-Leica,E</b>
Primary Accession	<a href="#">P25067</a>
Other Accession	<a href="#">NP_005193.1</a>
Reactivity	<b>Human, Mouse</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>Rabbit Ig</b>
Antigen Region	<b>571-599</b>

**COL8A2 Antibody (C-term) - Additional Information**

**Gene ID** 1296

**Other Names**

Collagen alpha-2(VIII) chain, Endothelial collagen, COL8A2

**Target/Specificity**

This COL8A2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 571-599 amino acids from the C-terminal region of human COL8A2.

**Dilution**

IF~~1:10~50  
WB~~1:2000  
IHC-P-Leica~~1:500


**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.


**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**



Confocal immunofluorescent analysis of COL8A2 Antibody (C-term)(Cat#AP11859b) with HepG2 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).



All lanes : Anti-COL8A2 Antibody (C-term) at 1:2000 dilution Lane 1: SH-SY5Y whole cell lysate Lane 2: Jurkat whole cell lysate Lane 3: Neuro-2a whole cell lysate Lane 4: C2C12 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG,

COL8A2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

#### COL8A2 Antibody (C-term) - Protein Information

**Name** COL8A2

#### Function

Macromolecular component of the subendothelium. Major component of the Descemet's membrane (basement membrane) of corneal endothelial cells. Also component of the endothelia of blood vessels. Necessary for migration and proliferation of vascular smooth muscle cells and thus, has a potential role in the maintenance of vessel wall integrity and structure, in particular in atherogenesis (By similarity).

#### Cellular Location

Secreted, extracellular space, extracellular matrix, basement membrane

#### Tissue Location


Expressed primarily in the subendothelium of large blood vessels. Also expressed in arterioles and venules in muscle, heart, kidney, spleen, umbilical cord, liver and lung and is also found in connective tissue layers around hair follicles, around nerve bundles in muscle, in the dura of the optic nerve, in cornea and sclera, and in the perichondrium of cartilaginous tissues. In the kidney, expressed in mesangial cells, glomerular endothelial cells, and tubular epithelial cells. Also expressed in mast cells, and in astrocytes during the repair process. Expressed in Descemet's membrane

#### COL8A2 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

(H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 67 kDa  
Blocking/Dilution buffer: 5% NFD/MTBST.



Immunohistochemical analysis of paraffin-embedded Human testis tissue using AP11859B performed on the Leica® BOND RXm. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:500) for 1 hours at room temperature. A undiluted biotinylated CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



Immunohistochemical analysis of paraffin-embedded Human kidney tissue using AP11859B performed on the Leica® BOND RXm. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA

- [Cell Culture](#)

buffer (pH9.0). Samples were incubated with primary antibody(1:500) for 1 hours at room temperature. A undiluted biotinylated CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.

### **COL8A2 Antibody (C-term) - Background**

This gene encodes the alpha 2 chain of type VIII collagen.

The protein is a major component of the basement membrane of the corneal endothelium and forms homo- or heterotrimers with alpha 1 (VIII) type collagens. Defects in this gene are associated with Fuchs endothelial corneal dystrophy and posterior polymorphous corneal dystrophy type 2.

### **COL8A2 Antibody (C-term) - References**

Hemadevi, B., et al. BMC Ophthalmol 10, 3 (2010) :  
Afshari, N.A., et al. Invest. Ophthalmol. Vis. Sci. 50(3):1093-1097(2009)  
Liskova, P., et al. Br J Ophthalmol 91(12):1717-1718(2007)  
Aldave, A.J., et al. Cornea 26(8):963-965(2007)  
Zhang, C., et al. Trans Am Ophthalmol Soc 104, 85-97 (2006) :

### **COL8A2 Antibody (C-term) - Citations**

- [Transcription factor TFAP2B up-regulates human corneal endothelial cell-specific genes during corneal development and maintenance.](#)
- [Involvement of ZEB1 and Snail1 in excessive production of extracellular matrix in Fuchs endothelial corneal dystrophy.](#)