

PPARD Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP11939b

Specification

PPARD Antibody (C-term) - Product Information

Application WB, FC,E **Primary Accession** Q03181 NP 803184.1 Other Accession Reactivity Human Host Rabbit Clonality **Polyclonal** Isotype Rabbit Ig Calculated MW 49903 Antigen Region 367-394

PPARD Antibody (C-term) - Additional Information

Gene ID 5467

Other Names

Peroxisome proliferator-activated receptor delta, PPAR-delta, NUCI, Nuclear hormone receptor 1, NUC1, Nuclear receptor subfamily 1 group C member 2, Peroxisome proliferator-activated receptor beta, PPAR-beta, PPARD, NR1C2, PPARB

Target/Specificity

This PPARD antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 367-394 amino acids from the C-terminal region of human PPARD.

Dilution


WB~~1:1000 FC~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw



PPARD Antibody (C-term) (Cat. #AP11939b) western blot analysis in HepG2 cell line lysates (35ug/lane). This demonstrates the PPARD antibody detected the PPARD protein (arrow).



PPARD Antibody (C-term) (Cat. #AP11939b) flow cytometric analysis of HepG2 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

PPARD Antibody (C-term) - Background



cycles.

Precautions

PPARD Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

PPARD Antibody (C-term) - Protein Information

Name PPARD

Synonyms NR1C2, PPARB

Function

Ligand-activated transcription factor.
Receptor that binds peroxisome
proliferators such as hypolipidemic drugs
and fatty acids. Has a preference for
poly-unsaturated fatty acids, such as
gamma- linoleic acid and eicosapentanoic
acid. Once activated by a ligand, the
receptor binds to promoter elements of
target genes. Regulates the peroxisomal
beta-oxidation pathway of fatty acids.
Functions as transcription activator for the
acyl-CoA oxidase gene. Decreases
expression of NPC1L1 once activated by a
ligand.

Cellular Location Nucleus.

Tissue Location

Ubiquitous with maximal levels in placenta and skeletal muscle

PPARD Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

This gene encodes a member of the peroxisome proliferator-activated receptor (PPAR) family. PPARs are nuclear hormone receptors that bind peroxisome proliferators and control the size and number of peroxisomes produced by cells. PPARs mediate a variety of biological processes, and may be involved in the development of several chronic diseases. including diabetes, obesity, atherosclerosis, and cancer. This protein is a potent inhibitor of ligand-induced transcription activity of PPAR alpha and PPAR gamma. It may function as an integrator of transcription repression and nuclear receptor signaling. The expression of this gene is found to be elevated in colorectal cancer cells. The elevated expression can be repressed by adenomatosis polyposis coli (APC), a tumor suppressor protein related to APC/beta-catenin signaling pathway. Knockout studies in mice suggested the role of this protein in myelination of the corpus callosum, lipid metabolism, and epidermal cell proliferation. Alternate splicing results in multiple transcript variants.

PPARD Antibody (C-term) - References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010) Christopoulos, P., et al. Ann. N. Y. Acad. Sci. 1205, 185-191 (2010): Dunn, S.E., et al. J. Exp. Med. 207(8):1599-1608(2010) Eynon, N., et al. Mitochondrion (2010) In press:

Jguirim-Souissi, I., et al. Genet. Mol. Res. 9(3):1326-1333(2010)