

STAR4 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12801a

Specification

STAR4 Antibody (N-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	O96DR4
Other Accession	NP_631903.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	23517
Antigen Region	8-37

STAR4 Antibody (N-term) - Additional Information

Gene ID 134429

Other Names

StAR-related lipid transfer protein 4, START domain-containing protein 4, StARD4, STAR4

Target/Specificity

This STAR4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 8-37 amino acids from the N-terminal region of human STAR4.

Dilution

WB~~1:1000
IHC-P~~1:10~50


Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.


Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions



STAR4 Antibody (N-term) (Cat. #AP12801a) western blot analysis in human placenta tissue lysates (35ug/lane). This demonstrates the STAR4 antibody detected the STAR4 protein (arrow).



STAR4 Antibody (N-term) (Cat. #AP12801a) immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of STAR4 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

STARD4 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

STARD4 Antibody (N-term) - Protein Information

Name STARD4

Function

Involved in the intracellular transport of cholesterol. Binds cholesterol or other sterols.

STARD4 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

STARD4 Antibody (N-term) - Background

Cholesterol homeostasis is regulated, at least in part, by sterol regulatory element (SRE)-binding proteins (e.g., SREBP1; MIM 184756) and by liver X receptors (e.g., LXRA; MIM 602423). Upon sterol depletion, LXRs are inactive and SREBPs are cleaved, after which they bind promoter SREs and activate genes involved in cholesterol biosynthesis and uptake. Sterol transport is mediated by vesicles or by soluble protein carriers, such as steroidogenic acute regulatory protein (STAR; MIM 600617). STAR is homologous to a family of proteins containing a 200- to 210-amino acid STAR-related lipid transfer (START) domain, including STARD4 (Soccio et al., 2002 [PubMed 12011452]).

STARD4 Antibody (N-term) - References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :
Korytowski, W., et al. Biochem. Biophys. Res. Commun. 392(1):58-62(2010)
Talmud, P.J., et al. Am. J. Hum. Genet. 85(5):628-642(2009)
Rodriguez-Agudo, D., et al. J. Lipid Res. 49(7):1409-1419(2008)