

ERO1L Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP13305b

Specification

ERO1L Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	O96HE7
Other Accession	NP_055399.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	54393
Antigen Region	350-379

ERO1L Antibody (C-term) - Additional Information

Gene ID 30001

Other Names

ERO1-like protein alpha, ERO1-L, ERO1-L-alpha, 184-, Endoplasmic oxidoreductin-1-like protein, Oxidoreductin-1-L-alpha, ERO1L

Target/Specificity

This ERO1L antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 350-379 amino acids from the C-terminal region of human ERO1L.

Dilution

WB~~1:1000
IHC-P~~1:10~50


Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.


Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions



ERO1L Antibody (C-term) (Cat. #AP13305b) western blot analysis in CEM cell line lysates (35ug/lane). This demonstrates the ERO1L antibody detected the ERO1L protein (arrow).



ERO1L Antibody (C-term) (Cat. #AP13305b) immunohistochemistry analysis in formalin fixed and paraffin embedded human stomach tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of ERO1L Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

ERO1L Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ERO1L Antibody (C-term) - Protein Information

Name ERO1A ([HGNC:13280](#))

Synonyms ERO1L

Function

Oxidoreductase involved in disulfide bond formation in the endoplasmic reticulum. Efficiently reoxidizes P4HB/PDI, the enzyme catalyzing protein disulfide formation, in order to allow P4HB to sustain additional rounds of disulfide formation. Following P4HB reoxidation, passes its electrons to molecular oxygen via FAD, leading to the production of reactive oxygen species (ROS) in the cell. Required for the proper folding of immunoglobulins. Involved in the release of the unfolded cholera toxin from reduced P4HB/PDI in case of infection by *V.cholerae*, thereby playing a role in retrotranslocation of the toxin. Plays an important role in ER stress-induced, CHOP-dependent apoptosis by activating the inositol 1,4,5-trisphosphate receptor IP3R1.

Cellular Location

Endoplasmic reticulum membrane; Peripheral membrane protein; Lumenal side. Note=The association with ERP44 is essential for its retention in the endoplasmic reticulum

Tissue Location

Widely expressed at low level. Expressed at high level in upper digestive tract. Highly expressed in esophagus. Weakly expressed in stomach and duodenum.

ERO1L Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

ERO1L Antibody (C-term) - Background

Essential oxidoreductase that oxidizes proteins in the endoplasmic reticulum to produce disulfide bonds. Acts by oxidizing directly P4HB/PDI isomerase through a direct disulfide exchange. Does not act as a direct oxidant of folding substrate, but relies on P4HB/PDI to transfer oxidizing equivalent. Associates with ERP44 but not with GRP54, demonstrating that it does not oxidize all PDI related proteins and can discriminate between PDI and related proteins. Its reoxidation probably involves electron transfer to molecular oxygen via FAD. Acts independently of glutathione. May be responsible for a significant proportion of reactive oxygen species (ROS) in the cell, thereby being a source of oxidative stress. Required for the folding of immunoglobulin proteins. Responsible for the release of the unfolded cholera toxin from reduced P4HB/PDI in case of infection by *V.cholerae*, thereby playing a role in retrotranslocation of the toxin.

ERO1L Antibody (C-term) - References

Inaba, K., et al. EMBO J. 29(19):3330-3343(2010)
Appenzeller-Herzog, C., et al. EMBO J. 29(19):3318-3329(2010)
Swiatkowska, M., et al. J. Biol. Chem. 285(39):29874-29883(2010)
Chambers, J.E., et al. J. Biol. Chem. 285(38):29200-29207(2010)
Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :

- [Flow Cytometry](#)
- [Cell Culture](#)