

MMP19 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP13759a

Specification

MMP19 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q99542
Other Accession	NP_002420.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	57357
Antigen Region	55-84

MMP19 Antibody (N-term) - Additional Information

Gene ID 4327

Other Names

Matrix metalloproteinase-19, MMP-19, 3424-, Matrix metalloproteinase RASI, Matrix metalloproteinase-18, MMP-18, MMP19, MMP18, RASI

Target/Specificity

This MMP19 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 55-84 amino acids from the N-terminal region of human MMP19.

Dilution

WB~~1:1000


Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions



MMP19 Antibody (N-term) - Background

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. This protein is expressed in human epidermis and it has a role in cellular proliferation as well as migration and adhesion to type I collagen. Multiple transcript

MMP19 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

MMP19 Antibody (N-term) - Protein Information

Name MMP19

Synonyms MMP18, RASI

Function

Endopeptidase that degrades various components of the extracellular matrix, such as aggrecan and cartilage oligomeric matrix protein (comp), during development, haemostasis and pathological conditions (arthritic disease). May also play a role in neovascularization or angiogenesis. Hydrolyzes collagen type IV, laminin, nidogen, nascin-C isoform, fibronectin, and type I gelatin.

Cellular Location

Secreted, extracellular space, extracellular matrix

Tissue Location

Expressed in mammary gland, placenta, lung, pancreas, ovary, small intestine, spleen, thymus, prostate, testis colon, heart and blood vessel walls. Not detected in brain and peripheral blood leukocytes. Also expressed in the synovial fluid of normal and rheumatoid patients (PubMed:8920941)

variants encoding distinct isoforms have been identified for this gene.

MMP19 Antibody (N-term) - References

Romero, R., et al. Am. J. Obstet. Gynecol. 203 (4), 361 (2010) :
Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Romero, R., et al. Am. J. Obstet. Gynecol. 202 (5), 431 (2010) :
Muller, M., et al. Mod. Pathol. 23(4):511-521(2010)
Lettau, I., et al. J. Neuropathol. Exp. Neurol. 69(3):215-223(2010)

MMP19 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)