

Mouse Cdk3 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP13803b

Specification

Mouse Cdk3 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q80YPO
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	33888
Antigen Region	249-277

Mouse Cdk3 Antibody (C-term) - Additional Information

Other Names

Cyclin-dependent kinase 3, Cell division protein kinase 3, Cdk3, Cdk3-ps, Cdkn3

Target/Specificity

This Mouse Cdk3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 249-277 amino acids from the C-terminal region of mouse Cdk3.

Dilution

WB~~1:1000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mouse Cdk3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.



Mouse Cdk3 Antibody (C-term) (Cat. #AP13803b) western blot analysis in CEM cell line lysates (35ug/lane). This demonstrates the Cdk3 antibody detected the Cdk3 protein (arrow).



Western blot analysis of Cdk3 (arrow) using rabbit polyclonal Mouse Cdk3 Antibody (C-term) (Cat. #AP13803b). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the Cdk3 gene.

Mouse Cdk3 Antibody (C-term) - Background

Probably involved in the control of the cell

Mouse Cdk3 Antibody (C-term) - Protein Information**Name** Cdk3**Synonyms** Cdk3-ps, Cdkn3**Function**

Serine/threonine-protein kinase that plays a critical role in the control of the eukaryotic cell cycle; involved in G0-G1 and G1-S cell cycle transitions. Interacts with CCNC/cyclin-C during interphase. Phosphorylates histone H1, ATF1, RB1 and CABLES1. ATF1 phosphorylation triggers ATF1 transactivation and transcriptional activities, and promotes cell proliferation and transformation. CDK3/cyclin-C mediated RB1 phosphorylation is required for G0-G1 transition. Promotes G1-S transition probably by contributing to the activation of E2F1, E2F2 and E2F3 in a RB1-independent manner.

cycle. Interacts with a yet unknown type of cyclin. Can phosphorylate histone H1 (By similarity).

Mouse Cdk3 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)