

USP16 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP16746a

Specification

USP16 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q9Y5T5
Other Accession	Q4R6X7 , NP_001001992.1 , NP_001027582.1
Reactivity	Human
Predicted	Monkey
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	93570
Antigen Region	132-160

USP16 Antibody (N-term) - Additional Information

Gene ID 10600

Other Names

Ubiquitin carboxyl-terminal hydrolase 16
{ECO:0000255|HAMAP-Rule:MF_03062},
341912
{ECO:0000255|HAMAP-Rule:MF_03062},
Deubiquitinating enzyme 16
{ECO:0000255|HAMAP-Rule:MF_03062},
Ubiquitin thioesterase 16
{ECO:0000255|HAMAP-Rule:MF_03062},
Ubiquitin-processing protease UBP-M,
Ubiquitin-specific-processing protease 16
{ECO:0000255|HAMAP-Rule:MF_03062},
USP16
{ECO:0000255|HAMAP-Rule:MF_03062}

Target/Specificity


This USP16 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 132-160 amino acids from the N-terminal region of human USP16.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This



USP16 Antibody (N-term) (Cat. #AP16746a) western blot analysis in SK-BR-3 cell line lysates (35ug/lane). This demonstrates the USP16 antibody detected the USP16 protein (arrow).

USP16 Antibody (N-term) - Background

This gene encodes a deubiquitinating enzyme that is phosphorylated at the onset of mitosis and then dephosphorylated at the metaphase/anaphase transition. It can deubiquitinate H2A, one of two major ubiquitinated proteins of chromatin, in vitro and a mutant form of the protein was shown to block cell division. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.

USP16 Antibody (N-term) - References

Gelsi-Boyer, V., et al. BMC Cancer 8, 299 (2008) :
Joo, H.Y., et al. Nature 449(7165):1068-1072(2007)
Pai, M.T., et al. J. Mol. Biol. 370(2):290-302(2007)
Sugiyama, N., et al. Mol. Cell Proteomics

antibody is purified through a protein A column, followed by peptide affinity purification.

6(6):1103-1109(2007)
Beausoleil, S.A., et al. Proc. Natl. Acad. Sci. U.S.A. 101(33):12130-12135(2004)

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

USP16 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

USP16 Antibody (N-term) - Protein Information

Name USP16

{ECO:0000255|HAMAP-Rule:MF_03062}

Function

Specifically deubiquitinates 'Lys-120' of histone H2A (H2AK119Ub), a specific tag for epigenetic transcriptional repression, thereby acting as a coactivator. Deubiquitination of histone H2A is a prerequisite for subsequent phosphorylation at 'Ser-11' of histone H3 (H3S10ph), and is required for chromosome segregation when cells enter into mitosis. In resting B- and T-lymphocytes, phosphorylation by AURKB leads to enhance its activity, thereby maintaining transcription in resting lymphocytes. Regulates Hox gene expression via histone H2A deubiquitination. Prefers nucleosomal substrates. Does not deubiquitinate histone H2B.

Cellular Location

Nucleus.

Tissue Location

Present in all the tissues examined including fetal brain, lung, liver, kidney, and adult heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas

USP16 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)