

METAP1 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17161c

Specification

METAP1 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	P53582
Other Accession	Q5ZIM5 , A6QLA4 , Q8BP48 , NP_055958.2
Reactivity Predicted	Human Mouse, Bovine, Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	43215
Antigen Region	188-216

METAP1 Antibody (Center) - Additional Information

Gene ID 23173

Other Names

Methionine aminopeptidase 1
{ECO:0000255|HAMAP-Rule:MF_03174},
MAP 1
{ECO:0000255|HAMAP-Rule:MF_03174},
MetAP 1
{ECO:0000255|HAMAP-Rule:MF_03174},
341118
{ECO:0000255|HAMAP-Rule:MF_03174},
Peptidase M 1
{ECO:0000255|HAMAP-Rule:MF_03174},
METAP1, KIAA0094

Target/Specificity


This METAP1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 188-216 amino acids from the Central region of human METAP1.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This



METAP1 Antibody (Center) - Background

METAP1 removes the amino-terminal methionine from nascent proteins. Required for normal progression through the cell cycle.

METAP1 Antibody (Center) - References

- Xiao, Q., et al. Biochemistry 49(26):5588-5599(2010)
- Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :
- Ross, C.J., et al. Nat. Genet. 41(12):1345-1349(2009)
- Hu, X.V., et al. Biochemistry 46(44):12833-12843(2007)
- Ewing, R.M., et al. Mol. Syst. Biol. 3, 89 (2007) :

antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

METAP1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

METAP1 Antibody (Center) - Protein Information

Name METAP1

Synonyms KIAA0094

Function

Cotranslationally removes the N-terminal methionine from nascent proteins. The N-terminal methionine is often cleaved when the second residue in the primary sequence is small and uncharged (Met- Ala-, Cys, Gly, Pro, Ser, Thr, or Val). Required for normal progression through the cell cycle.

Cellular Location

Cytoplasm
{ECO:0000255|HAMAP-Rule:MF_03174}.

METAP1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)