

MYRIP Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17165B

Specification

MYRIP Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q8NFW9
Other Accession	NP_056275.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	95706
Antigen Region	659-686

MYRIP Antibody (C-term) - Additional Information

Gene ID 25924

Other Names

Rab effector MyRIP, Exophilin-8, Myosin-VIIa- and Rab-interacting protein, Synaptotagmin-like protein lacking C2 domains C, SlaC2-c, Slp homolog lacking C2 domains c, MYRIP, SLAC2C

Target/Specificity

This MYRIP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 659-686 amino acids from the C-terminal region of human MYRIP.

Dilution

WB~~1:1000


Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions



MYRIP Antibody (C-term) - Background

Rab effector protein involved in melanosome transport. Serves as link between melanosome-bound RAB27A and the motor proteins MYO5A and MYO7A. May link RAB27A-containing vesicles to actin filaments (By similarity).

MYRIP Antibody (C-term) - References

Rose, J. Phd, et al. Mol. Med. (2010) In press :
Nightingale, T.D., et al. Blood 113(20):5010-5018(2009)
Hayasaka, S., et al. Asian J. Androl. 10(4):561-568(2008)
Goehring, A.S., et al. J. Biol. Chem. 282(45):33155-33167(2007)
Wu, C., et al. Proteomics 7(11):1775-1785(2007)

MYRIP Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

MYRIP Antibody (C-term) - Protein Information

Name MYRIP

Synonyms SLAC2C

Function

Rab effector protein involved in melanosome transport. Serves as link between melanosome-bound RAB27A and the motor proteins MYO5A and MYO7A. May link RAB27A-containing vesicles to actin filaments. Functions as a protein kinase A-anchoring protein (AKAP). May act as a scaffolding protein that links PKA to components of the exocytosis machinery, thus facilitating exocytosis, including insulin release (By similarity).

Cellular Location

Cytoplasm

{ECO:0000250|UniProtKB:Q8K3I4}.

Cytoplasm, perinuclear region

{ECO:0000250|UniProtKB:Q7TNY7}

Cytoplasmic vesicle, secretory vesicle

{ECO:0000250|UniProtKB:Q7TNY7}

Note=In presynaptic and postsynaptic areas in photoreceptor cells and in the basal microvilli of retinal pigment epithelium cells. Associated with melanosomes. Colocalizes with actin filaments

{ECO:0000250|UniProtKB:Q7TNY7}

Tissue Location

Detected in brain, skin, heart, adrenal medulla, pancreas, intestine, liver, kidney, muscle and testis

MYRIP Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)