

EIF3E Antibody(Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP19409c

Specification

EIF3E Antibody(Center) - Product Information

| | |
|----------------------|---|
| Application | WB,E |
| Primary Accession | P60228 |
| Other Accession | Q641X8 , P60229 , Q4R6G8 , Q5ZLA5 , Q3T102 , Q3B8M3 , Q1LUA8 , Q05AY2 , Q6DRI1 , NP_001559.1 |
| Reactivity Predicted | Human, Zebrafish Xenopus, Bovine, Chicken, Monkey, Mouse, Rat |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit Ig |
| Calculated MW | 52221 |
| Antigen Region | 248-276 |

EIF3E Antibody(Center) - Additional Information

Gene ID 3646


Other Names

Eukaryotic translation initiation factor 3 subunit E
 {ECO:0000255|HAMAP-Rule:MF_03004},
 eIF3e
 {ECO:0000255|HAMAP-Rule:MF_03004},
 Eukaryotic translation initiation factor 3 subunit 6
 {ECO:0000255|HAMAP-Rule:MF_03004},
 Viral integration site protein INT-6 homolog,
 eIF-3 p48
 {ECO:0000255|HAMAP-Rule:MF_03004},
 EIF3E
 {ECO:0000255|HAMAP-Rule:MF_03004}


Target/Specificity

This EIF3E antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 248-276 amino acids from the Central region of human EIF3E.

Dilution



(DANRE) eif3eb Antibody (Center) (Cat.# AP19409c) western blot analysis in zebra fish brain tissue lysates (35ug/lane). This demonstrates the (DANRE) eif3eb antibody detected the (DANRE) eif3eb protein (arrow).



EIF3E Antibody (Center)(Cat. #AP19409c) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the EIF3E antibody detected the EIF3E protein (arrow).

EIF3E Antibody(Center) - Background

Component of the eukaryotic translation

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

EIF3E Antibody(Center) is for research use only and not for use in diagnostic or therapeutic procedures.

EIF3E Antibody(Center) - Protein Information

Name EIF3E

{ECO:0000255|HAMAP-Rule:MF_03004}

Function

Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis (PubMed:17581632, PubMed:25849773, PubMed:27462815). The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl- tRNA_i and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation (PubMed:17581632). The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling,

initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNA_i and eIF-5 to form the 43S preinitiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of posttermination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. Required for nonsense-mediated mRNA decay (NMD); may act in conjunction with UPF2 to divert mRNAs from translation to the NMD pathway. May interact with MCM7 and EPAS1 and regulate the proteasome-mediated degradation of these proteins.

EIF3E Antibody(Center) - References

- Grzmil, M., et al. *Oncogene* 29(28):4080-4089(2010)
Zhou, M., et al. *Proc. Natl. Acad. Sci. U.S.A.* 105(47):18139-18144(2008)
Masutani, M., et al. *EMBO J.* 26(14):3373-3383(2007)
Morris, C., et al. *EMBO Rep.* 8(6):596-602(2007)
Sirchia, R., et al. *Biol. Chem.* 388(5):457-465(2007)

differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression (PubMed:[25849773](http://www.uniprot.org/citations/25849773)). Required for nonsense-mediated mRNA decay (NMD); may act in conjunction with UPF2 to divert mRNAs from translation to the NMD pathway (PubMed:[17468741](http://www.uniprot.org/citations/17468741)). May interact with MCM7 and EPAS1 and regulate the proteasome-mediated degradation of these proteins (PubMed:[17310990](http://www.uniprot.org/citations/17310990), PubMed:[17324924](http://www.uniprot.org/citations/17324924)).

Cellular Location

Cytoplasm. Nucleus, PML body.

Tissue Location

Ubiquitously expressed. Expressed at highest levels in appendix, lymph, pancreas, skeletal muscle, spleen and thymus

EIF3E Antibody(Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)