

CTNS Antibody(C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP19410b

Specification

CTNS Antibody(C-term) - Product Information

Application	WB,E
Primary Accession	O60931
Other Accession	NP_004928.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	341-367

CTNS Antibody(C-term) - Additional Information

Gene ID 1497

Other Names
Cystinosis, CTNS

Target/Specificity

This CTNS antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 341-367 amino acids from the C-terminal region of human CTNS.

Dilution

WB~~1:1000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CTNS Antibody(C-term) is for research use only and not for use in diagnostic or therapeutic procedures.



CTNS Antibody(C-term) - Background

This gene encodes a seven-transmembrane domain protein that functions to transport cystine out of lysosomes. Its activity is driven by the H⁺ electrochemical gradient of the lysosomal membrane. Mutations in this gene cause cystinosis, a lysosomal storage disorder. Alternative splicing results in multiple transcript variants.

CTNS Antibody(C-term) - References

- Taranta, A., et al. *Pediatr. Nephrol.* 25(7):1263-1267(2010)
- Bellomo, F., et al. *Free Radic. Biol. Med.* 48(7):865-872(2010)
- Kumar, A., et al. *J. Biosci.* 35(1):21-25(2010)
- Aldahmesh, M.A., et al. *Ophthalmic Genet.* 30(4):185-189(2009)
- Macias-Vidal, J., et al. *Clin. Genet.* 76(5):486-489(2009)

CTNS Antibody(C-term) - Protein Information**Name** CTNS

{ECO:0000303|PubMed:9537412,
ECO:0000312|HGNC:HGNC:2518}

Function

Cystine/H(+) symporter that mediates export of cystine, the oxidized dimer of cysteine, from lysosomes (PubMed:11689434, PubMed:18337546, PubMed:22232659, PubMed:29467429, PubMed:33208952, PubMed:15128704). Plays an important role in melanin synthesis by catalyzing cystine export from melanosomes, possibly by inhibiting pheomelanin synthesis (PubMed:22649030). In addition to cystine export, also acts as a positive regulator of mTORC1 signaling in kidney proximal tubular cells, via interactions with components of the v-ATPase and Ragulator complexes (By similarity). Also involved in small GTPase-regulated vesicle trafficking and lysosomal localization of LAMP2A, independently of cystine transporter activity (By similarity).

Cellular Location

[Isoform 1]: Lysosome membrane; Multi-pass membrane protein. Melanosome membrane; Multi-pass membrane protein. Note=AP-3 complex is required for localization to the lysosome.

Tissue Location

Strongly expressed in pancreas, kidney (adult and fetal), skeletal muscle, melanocytes and keratinocytes (PubMed:22649030). Expressed at lower levels in placenta and heart Weakly

expressed in lung, liver and brain (adult and fetal) (PubMed:22649030).

CTNS Antibody(C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)