

### **COPS5 Antibody (Center)**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP21563c

## **Specification**

**COPS5 Antibody (Center) - Product Information** 

Application	WB,E
Primary Accession	<u>Q92905</u>
Reactivity	Human, Mouse,
	Rat
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit Ig
Calculated MW	37579

#### **COPS5 Antibody (Center) - Additional Information**

#### Gene ID 10987

#### **Other Names**

COP9 signalosome complex subunit 5, SGN5, Signalosome subunit 5, 34--, Jun activation domain-binding protein 1, COPS5, CSN5, JAB1

### Target/Specificity

This COPS5 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 187-222 amino acids from the Central region of human COPS5.

**Dilution** WB~~1:500-1:1000

### Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

### Precautions

COPS5 Antibody (Center) is for research use only and not for use in diagnostic or



All lanes : Anti-COPS5 Antibody (Center) at 1:500-1:1000 dilution Lane 1: NIH/3T3 whole cell lysates Lane 2: U-87 MG whole cell lysates Lane 3: PC-12 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 38 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

# **COPS5 Antibody (Center) - Background**

Probable protease subunit of the COP9 signalosome complex (CSN), a complex involved in various cellular and developmental processes. The CSN complex is an essential regulator of the ubiquitin (Ubl) conjugation pathway by mediating the deneddylation of the cullin subunits of the SCF-type E3 ligase complexes, leading to decrease the Ubl ligase activity of SCF-type complexes such as SCF, CSA or DDB2. The complex is also involved in phosphorylation of p53/TP53, c-jun/JUN, IkappaBalpha/NFKBIA, ITPK1 and IRF8, possibly via its association with CK2 and PKD kinases. CSN-dependent phosphorylation of TP53 and JUN promotes and protects degradation by the Ubl system, respectively. In the complex, it



therapeutic procedures.

**COPS5 Antibody (Center) - Protein Information** 

Name COPS5

Synonyms CSN5, JAB1

## Function

Probable protease subunit of the COP9 signalosome complex (CSN), a complex involved in various cellular and developmental processes. The CSN complex is an essential regulator of the ubiquitin (Ubl) conjugation pathway by mediating the deneddylation of the cullin subunits of the SCF-type E3 ligase complexes, leading to decrease the Ubl ligase activity of SCF-type complexes such as SCF, CSA or DDB2. The complex is also involved in phosphorylation of p53/TP53, c-jun/JUN,

IkappaBalpha/NFKBIA, ITPK1 and IRF8, possibly via its association with CK2 and PKD kinases. CSN-dependent phosphorylation of TP53 and JUN promotes and protects degradation by the Ubl system, respectively. In the complex, it probably acts as the catalytic center that mediates the cleavage of Nedd8 from cullins. It however has no metalloprotease activity by itself and requires the other subunits of the CSN complex. Interacts directly with a large number of proteins that are regulated by the CSN complex. confirming a key role in the complex. Promotes the proteasomal degradation of BRSK2.

### **Cellular Location**

Cytoplasm, cytosol. Nucleus. Cytoplasm, perinuclear region. Cytoplasmic vesicle, secretory vesicle, synaptic vesicle Note=Nuclear localization is diminished in the presence of IFIT3

# **COPS5 Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- <u>Dot Blot</u>
- Immunohistochemistry

probably acts as the catalytic center that mediates the cleavage of Nedd8 from cullins. It however has no metalloprotease activity by itself and requires the other subunits of the CSN complex. Interacts directly with a large number of proteins that are regulated by the CSN complex, confirming a key role in the complex. Promotes the proteasomal degradation of BRSK2.

# **COPS5** Antibody (Center) - References

Claret F.-X.,et al.Nature 383:453-457(1996). Asano K.,et al.J. Biol. Chem. 272:27042-27052(1997). Halleck A.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases. Bechtel S.,et al.BMC Genomics 8:399-399(2007). Bienvenut W.V.,et al.Submitted (OCT-2005) to UniProtKB.



- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
  Cell Culture