

# Phospho-IL6ST(Y905) Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP3663a

## **Specification**

# Phospho-IL6ST(Y905) Antibody - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Isotype
Calculated MW

DB,E
P40189
Human
Rabbit
Polyclonal
Rabbit Ig
103537

Phospho-IL6ST(Y905) Antibody - Additional Information

#### Gene ID 3572

#### Other Names

Interleukin-6 receptor subunit beta, IL-6 receptor subunit beta, IL-6R subunit beta, IL-6R-beta, IL-6RB, CDw130, Interleukin-6 signal transducer, Membrane glycoprotein 130, gp130, Oncostatin-M receptor subunit alpha, CD130, IL6ST

### **Target/Specificity**

This IL6ST Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding Y905 of human IL6ST.

#### Dilution


DB~~1:500

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

### **Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.




Dot blot analysis of anti-Phospho-IL6ST-Y905 Phospho-specific Pab (Cat. #AP3663a) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per ml.

# Phospho-IL6ST(Y905) Antibody - Background

IL6ST is a signal transducer shared by many cytokines, including interleukin 6 (IL6), ciliary neurotrophic factor (CNTF), leukemia inhibitory factor (LIF), and oncostatin M (OSM). This protein functions as a part of the cytokine receptor complex. The activation of this protein is dependent upon the binding of cytokines to their receptors. vIL6, a protein related to IL6 and encoded by the Kaposi sarcoma-associated herpesvirus, can bypass the interleukin 6 receptor (IL6R) and directly activate this protein.

# Phospho-IL6ST(Y905) Antibody - References

Lehmann, U., et.al., J. Biol. Chem. 278 (1), 661-671 (2003)





**Precautions** 

Phospho-IL6ST(Y905) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.


Phospho-IL6ST(Y905) Antibody - Protein Information

Name IL6ST (HGNC:6021)

#### **Function**

Signal-transducing molecule (PubMed:<a hr ef="http://www.uniprot.org/citations/22616 37" target=" blank">2261637</a>). The receptor systems for IL6, LIF, OSM, CNTF, IL11, CTF1 and BSF3 can utilize IL6ST for initiating signal transmission. Binding of IL6 to IL6R induces IL6ST homodimerization and formation of a high-affinity receptor complex, which activate the intracellular JAK-MAPK and JAK-STAT3 signaling pathways (PubMed: <a href="http://www.uni prot.org/citations/2261637" target="\_blank">2261637</a>, PubMed:<a href="http://www.uniprot.org/ci tations/19915009" target=" blank">19915009</a>, PubMed:<a href="http://www.uniprot.org/ci tations/23294003" target="\_blank">23294003</a>). That causes phosphorylation of IL6ST tyrosine residues which in turn activates STAT3 (PubMed:<a href="http://www.uniprot.org/c itations/19915009" target=" blank">19915009</a>, PubMed:<a href="http://www.uniprot.org/ci tations/23294003" target="\_blank">23294003</a>, PubMed:<a href="http://www.uniprot.org/ci tations/25731159" target=" blank">25731159</a>). In parallel, the IL6 signaling pathway induces the expression of two cytokine receptor signaling inhibitors, SOCS1 and SOCS3, which inhibit JAK and terminate the activity of the IL6 signaling pathway as a negative feedback loop (By similarity). Also activates the yes- associated protein 1 (YAP) and NOTCH pathways to control inflammationinduced epithelial regeneration, independently of STAT3 (By similarity). Mediates signals which regulate immune response, hematopoiesis, pain control and bone metabolism (By similarity). Has a role in embryonic development (By similarity). Essential for survival of motor and sensory

Novotny-Diermayr, V., et.al., J. Biol. Chem. 277 (51), 49134-49142 (2002)





neurons and for differentiation of astrocytes (By similarity). Required for expression of TRPA1 in nociceptive neurons (By similarity). Required for the maintenance of PTH1R expression in the osteoblast lineage and for the stimulation of PTH-induced osteoblast differentiation (By similarity). Required for normal trabecular bone mass and cortical bone composition (By similarity).

#### **Cellular Location**

[Isoform 1]: Cell membrane; Single-pass type I membrane protein

# **Tissue Location**

Found in all the tissues and cell lines examined (PubMed:2261637). Expression not restricted to IL6 responsive cells (PubMed:2261637).

# Phospho-IL6ST(Y905) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture