

**Phospho-rat TSC1(Y297) Antibody**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP3832a**

## Specification

### Phospho-rat TSC1(Y297) Antibody - Product Information

Application	DB,E
Primary Accession	<a href="#">Q9Z136</a>
Other Accession	<a href="#">NP_068626.1</a>
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	129022

### Phospho-rat TSC1(Y297) Antibody - Additional Information

Gene ID 60445

### Other Names

Hamartin, Tuberous sclerosis 1 protein homolog, Tsc1

### Target/Specificity

This rat TSC1 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding Y297 of rat TSC1.

### Dilution

DB~~1:500

### Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

### Precautions

Phospho-rat TSC1(Y297) Antibody is for research use only and not for use in



### Phospho-rat TSC1(Y297) Antibody - Background

In complex with TSC2, inhibits the nutrient-mediated or growth factor-stimulated phosphorylation of S6K1 and EIF4EBP1 by negatively regulating mTORC1 signaling (By similarity). Implicated as a tumor suppressor. Involved in microtubule-mediated protein transport, but this seems to be due to unregulated mTOR signaling (By similarity).

### Phospho-rat TSC1(Y297) Antibody - References

- Inoue, H., et al. Biosci. Biotechnol. Biochem. 73(11):2488-2493(2009)  
Di Nardo, A., et al. J. Neurosci. 29(18):5926-5937(2009)  
Chen, P., et al. Exp. Mol. Pathol. 86(2):101-107(2009)

diagnostic or therapeutic procedures.

Momose, S., et al. Biochem. Biophys. Res. Commun. 356(3):693-698(2007)  
Goncharova, E., et al. J. Cell Biol. 167(6):1171-1182(2004)

#### **Phospho-rat TSC1(Y297) Antibody - Protein Information**

#### **Name Tsc1**

#### **Function**

In complex with TSC2, inhibits the nutrient-mediated or growth factor-stimulated phosphorylation of S6K1 and EIF4EBP1 by negatively regulating mTORC1 signaling (By similarity). Implicated as a tumor suppressor. Involved in microtubule-mediated protein transport, but this seems to be due to unregulated mTOR signaling (PubMed:<a href="http://www.uniprot.org/citations/16707451" target="\_blank">16707451</a>). Acts as a co-chaperone for HSP90AA1 facilitating HSP90AA1 chaperoning of protein clients such as kinases, TSC2 and glucocorticoid receptor NR3C1 (By similarity). Increases ATP binding to HSP90AA1 and inhibits HSP90AA1 ATPase activity (By similarity). Competes with the activating co-chaperone AHSA1 for binding to HSP90AA1, thereby providing a reciprocal regulatory mechanism for chaperoning of client proteins (By similarity). Recruits TSC2 to HSP90AA1 and stabilizes TSC2 by preventing the interaction between TSC2 and ubiquitin ligase HERC1 (By similarity).

#### **Cellular Location**

Cytoplasm  
{ECO:0000250|UniProtKB:Q92574}.  
Membrane  
{ECO:0000250|UniProtKB:Q92574};  
Peripheral membrane protein  
{ECO:0000250|UniProtKB:Q92574}.  
Note=At steady state found in association with membranes.  
{ECO:0000250|UniProtKB:Q92574}

#### **Tissue Location**

Highly expressed in brain, spleen and kidney, followed by liver and heart

#### **Phospho-rat TSC1(Y297) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)