

Anti-GANP Antibody

Catalog # AP53715

Specification

Anti-GANP Antibody - Product Information

Application WB, IH Primary Accession 060318

Reactivity Human, Mouse,

Rat

Host Rabbit
Clonality Polyclonal
Calculated MW 218405

Anti-GANP Antibody - Additional Information

Gene ID 8888

Other Names

GANP; KIAA0572; MAP80; Germinal-center associated nuclear protein; GANP; 80 kDa MCM3-associated protein; MCM3 acetylating protein; MCM3AP; MCM3 acetyltransferase

Target/Specificity

Recognizes endogenous levels of GANP protein.

Dilution

WB~~1/500 - 1/1000 IH~~1/50 - 1/200

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

Storage

Store at -20 °C.Stable for 12 months from date of receipt


Anti-GANP Antibody - Protein Information

Name MCM3AP


Synonyms GANP, KIAA0572, MAP80

Function

[Isoform GANP]: As a component of the TREX-2 complex, involved in the export of




Western blot analysis of GANP expression in rat testis (A), CT26 (B), SKOVCAR3 (C), HCT116 (D), HEK293T (E), HCC827 (F) whole cell lysates.



Immunohistochemical analysis of GANP staining in human tonsil formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Anti-GANP Antibody - Background





mRNAs to the cytoplasm through the nuclear pores (PubMed:20005110, PubMed:20384700

target="_blank">20384790,

PubMed:<a href="http://www.uniprot.org/ci tations/23591820"

target="_blank">23591820,

PubMed:<a href="http://www.uniprot.org/ci tations/22307388"

target="_blank">22307388). Through the acetylation of histones, affects the assembly of nucleosomes at immunoglobulin variable region genes and promotes the recruitment and positioning of transcription complex to favor DNA cytosine deaminase AICDA/AID targeting, hence promoting somatic hypermutations (PubMed:<a href="http://www.uniprot.org/citations/23652018"

itations/23652018" target=" blank">23652018).

Cellular Location

[Isoform GANP]: Nucleus envelope. Nucleus, nuclear pore complex. Nucleus, nucleoplasm. Chromosome.

Note=Predominantly located at the nuclear envelope, facing the nucleus interior (PubMed:20005110, PubMed:21195085, PubMed:23591820). Localization at the nuclear pore complex requires NUP153, TPR and ALYREF/ALY (PubMed:23591820, PubMed:22307388). Also found associated with chromatin (PubMed:23652018). In B-cells, targeted to the immunoglobulin variable region genes (PubMed:23652018).

Tissue Location


Widely expressed (PubMed:11024281). Up-regulated in germinal center B-cells in tonsils (at protein level) (PubMed:11024281).

Anti-GANP Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence

Rabbit polyclonal antibody to GANP





ImmunoprecipitationFlow CytometyCell Culture