

Anti-ZNF668 Antibody

Catalog # AP53913

Specification

Anti-ZNF668 Antibody - Product Information

Application	WB, IH, IF
Primary Accession	Q96K58
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	67890

Anti-ZNF668 Antibody - Additional Information

Gene ID 79759

Other Names

Zinc finger protein 668

Target/Specificity

Recognizes endogenous levels of ZNF668 protein.

Dilution

WB~~1/500 - 1/1000
IH~~1/50 - 1/200
IF~~1/50 - 1/200

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-ZNF668 Antibody - Protein Information


Name ZNF668

Function


May be involved in transcriptional regulation.

Cellular Location

Nucleus.



Western blot analysis of ZNF668 expression in mouse lung (A), mouse kidney (B), rat kidney (C) whole cell lysates.



Immunohistochemical analysis of ZNF668 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Anti-ZNF668 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



Immunofluorescent analysis of ZNF668 staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Anti-ZNF668 Antibody - Background

Rabbit polyclonal antibody to ZNF668