

FABP12 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55051

Specification

FABP12 Polyclonal Antibody - Product Information

Application IHC-P Primary Accession A6NFH5

Reactivity Rat, Dog, Cow

Host Rabbit
Clonality Polyclonal
Calculated MW 15565

FABP12 Polyclonal Antibody - Additional Information

Gene ID 646486

Other Names

Fatty acid-binding protein 12, FABP12

Format

0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

FABP12 Polyclonal Antibody - Protein Information

Name FABP12

Function


May play a role in lipid transport.

Tissue Location


Expressed in a number of retinoblastoma cell lines.

FABP12 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.



Paraformaldehyde-fixed, paraffin embedded (Human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FABP12) Polyclonal Antibody, Unconjugated (bs-13064R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FABP12) Polyclonal Antibody, Unconjugated





- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

(bs-13064R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.